



# Study of Subacute Toxicity in Wistar Rats Challenged with *Phyllanthus amarus* Schum and Thonn

Kpomah Enyohwo Dennis <sup>a\*</sup>, Kpomah Bridget <sup>b</sup>  
and Okonkwo Chisom Stanislaus <sup>c</sup>

<sup>a</sup> Biochemistry Department, Federal University Otuoke, Bayelsa State, Nigeria.

<sup>b</sup> Chemistry Department, Delta State University, Abraka, Nigeria.

<sup>c</sup> Science Laboratory Technology Department, Federal Polytechnic Oko, Anambra State, Nigeria.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jocamr/2024/v25i8558>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119874>

Original Research Article

Received: 16/05/2024

Accepted: 18/07/2024

Published: 05/08/2024

## ABSTRACT

*Phyllanthus amarus*, a widely used plant in complementary and alternative medicine practice in Southern Nigeria, is used in treating and managing numerous metabolic disorders, and neurodegenerative diseases. This study is designed to assess the subacute toxicity of *P. amarus* in Wistar rats using body and relative organ weight, renal function, *in-vivo* antioxidant status and organ histology indices as a baseline. The twenty adult male rats weighing  $120.00 \pm 6.18$  g were apportioned into four groups with five rats per group. Group A (Control) received 1.0 mL of distilled water, group B received 1000 mg/kg group C received 1500 mg/kg and group D received 2000

\*Corresponding author: E-mail: [kpomahed@fuotuo.ke.edu.ng](mailto:kpomahed@fuotuo.ke.edu.ng);

mg/kg body weight of the crude ethanol extract. The extract from *P. amarus* was administered orally once daily at 8:30 am using an oral cannula attached to a 2 mL syringe. Subacute toxicity was evaluated after 14 days. The findings showed no visible and noticeable overt signs of toxicity throughout the experimental period, non-significant ( $p > 0.05$ ) adverse change in body and relative organ weight, renal function and organ histopathology of the rats in the treated and control groups. However, *P. amarus* significantly ( $p < 0.05$ ) improved *in-vivo* antioxidant status while significantly reducing ( $p < 0.05$ ) the level of malondialdehyde, a biological indicator of oxidative stress in the living system.

**Keywords:** Antioxidants; oxidative stress; *Phyllanthus amarus*; phytochemicals.

## 1. INTRODUCTION

Medicines from plant sources commonly referred to as herbal medicine are an important alternative therapeutic aid for both curative and prophylactic purposes in developing and developed countries, primarily because of their wide range and diversity of phytochemicals inherent in them and their use dates back to antiquity. These phytochemicals include the different classes of flavonoids, tannins, saponins, and glycosides. Herbal medicine often involves using one or more plant components such as leaves, flowers, stem (bark) and roots, or a combination of two or more plants that work synergistically to heighten therapeutic potentials and other benefits. The active ingredients in these plant materials for medicinal purposes, if adhered to strictly in dose and regimen, are safe and reliable [1,2], cheap and nearly often available all year round [3]. Drugs from plants have been used with proven track records in the treatment and management of malaria, diabetes, hypertension, infertility, erectile dysfunction and management of hypoactive sexual disorders in men and women, cardiovascular disease, neurological diseases etc. [4,5].

*Phyllanthus amarus* is one such plant frequently used in tropical and subtropical countries with ethnomedicinal applications as antimicrobial, anti-inflammatory, antidiabetic, anticancer, and antiplasmodial [6]. *P. amarus* also possesses potent antioxidant and diuretic properties [7]. "*P. amarus*, a member of the family *Euphorbiaceae* and the genus *Phyllanthus* and the species is *amarus*. it has nearly 800 species and is widely distributed in tropical and subtropical countries" [8]. "There is a general belief by herbal practitioners and users of herbal medicine that because herbal products are sourced from nature they are therefore free of adverse or toxic effects unlike most synthetic drugs" [9,10]. Hence, the toxicity and adverse effects of most herbal products are often not evaluated and as

such the users often look at the therapeutic advantage of the plant and disregard their toxic effects on the body's vital organs and tissues.

Arising from the widespread usage of *P. amarus* in folklore medicine, this study seeks to appraise the sub-acute toxicity of *P. amarus* for safety or possible toxic effects using alterations in body and organ weight, antioxidant status, renal function activities and liver histology as indices of toxicity in rats.

## 2. MATERIALS AND METHODS

### 2.1 Sample Collection and Identification

*P. amarus* was harvested around the Faculty Building (FB1) Laboratory of the Federal University Otuoke, Bayelsa State, Nigeria on the 15<sup>th</sup> day of March 2024. The Plant Science section of the Biology Department, Federal University Otuoke, Bayelsa State identified and confirmed the plant sample.

### 2.2 Preparation of Plant Extract

The samples were painstakingly washed with distilled water to eliminate trash and contaminants, it was then air dried for 14 days to give a persistent weight and then milled using an electric blender (Blender 462 Nakai Japan). 100 g of the powdered *P. amarus* was extracted in 300 mL of absolute ethanol for 24 hours at room temperature with continuous shaking using a flask shaker (Denly A 500). The extract was filtered with WhatmanNo.1 filter paper and the resultant filtrate evaporated to dryness using a rotary evaporator at 40°C to give 4.33 g of the crude extract.

### 2.3 Experimental Animals

Twenty healthy, male albino rats weighing 120.00 ± 6.18 g were purchased from the Animal

House section, Department of Biochemistry, Federal University Otuoke, Bayelsa State, Nigeria. The animals were kept in separate investigational rooms, which were clean and well-ventilated at a temperature between 28-30°C, under a natural dark/light cycle with free access to standard rat chow and water *ad-libitum* during the period of acclimatization which lasted for one week.

## 2.4 Animal Grouping

The twenty adult male rats were divided into four groups with five rats per group. Group A (Control), received 1.0 mL of distilled water, group B, received 1000 mg/kg, group C, received 1500 mg/kg and group D, received 2000 mg/kg body weight of the extract. The extract from *P. amarus* was administered orally once daily at 8:30 am using an oral cannula attached to a 2 mL syringe. These doses were carefully chosen to avoid the LD<sub>50</sub> but marginally above the effective dose of *P. amarus*. The rats were fed *ad libitum* with standard chow and tap water throughout the experimental protocol. The study was carried out for 14 days and on the 15th day, the animals were sacrificed.

## 2.5 Weekly Cage Side Surveillance for Physical Signs of Toxicity

This was done by physical examination of the rats from the cage sides for overt signs of toxicity such as salivation, lacrimation, eye dullness, eye opacity, diarrhoea, restlessness, red stained muzzle, lethargy, piloerection, skin appearance, subcutaneous swelling, loss of appetite, colour and consistency of faeces abdominal distension and mortality arising from the administration of the different doses of *P. amarus*.

### 2.5.1 Changes in body weight

Rats in all groups were weighed on the first day and after the treatment protocols. Percentage change in body weight was evaluated by the expression below [11]:

$$\begin{aligned} & \% \text{ change in body weight} \\ & = \frac{\text{final body weight} - \text{initial body weight}}{\text{initial body weight}} \times 100\% \end{aligned}$$

### 2.5.2 Organ (kidney, liver and heart) as the ratio of body weight

Organs (kidney and liver) were removed and weighed immediately. The organ ratio was

evaluated as a percentage with the expression below [11].

$$\begin{aligned} & \text{Organ ratio (\%)} \\ & = \frac{\text{weight of organ (g)}}{\text{body weight (g)}} \times 100\% \end{aligned}$$

## 2.6 Animal Handling Procedure and Sample Preparation Technique

On day 15<sup>th</sup> of the experimental protocol, the rats were euthanized under anesthesia using diethyl ether chamber, blood specimen were obtained by cardiac puncture into plain sample bottles. The blood specimen was allowed to stand for 20 minutes for coagulation to occur, afterwards, the specimen was centrifuged at 2000 rpm for 10 minutes and the supernatant (serum) was collected and stored in the refrigerator before the biochemical assay. The liver and kidney were dissected instantaneously for histological studies.

## 2.7 Biochemical Assay Kits

Assay kits for renal function indices and antioxidants are products of Randox Laboratories Ltd., United Kingdom. All other reagents/chemicals were obtained from standard suppliers and of analytical grade.

### 2.7.1 Biochemical analysis of antioxidant enzymes

Catalase activity was estimated by the method of Cohen et al. [12]. "Superoxide dismutase (SOD) activity was by the methods of Misra and Fridovich" [13]. The assay method of Hunter et al. [14] as modified by Gutteridge and Wilkins [15] was adopted for the assay of Malondialdehyde (MDA) concentration.

### 2.7.2 Biochemical analysis of renal function

Blood urea nitrogen (BUN) was evaluated by the Berthelot method as modified by Tobacco et al. [16]. Creatinine (CRT) was assayed by the colourimetric kinetic method of Bartels et al. [17]. Uric acid (UA) was assessed using the enzymatic colourimetric method of Duncan et al. [18].

## 2.8 Histopathological Scrutiny of the Organs

"The liver and kidney slices for histopathology were fixed in 10% formal saline and embedded in

paraffin wax blocks, sections of 5µm thick were stained with hematoxylin and eosin (H&E) and then examined under a light microscope for determination of derangement and pathological changes” [19,20].

## 2.9 Statistical Analysis

Experimental values were expressed as means ± SD. To determine differences between the groups studied, a one-way analysis of variance (ANOVA) with Duncan post hoc test was used to compare the group means, and  $p < 0.05$  was considered statistically significant. SPSS for Windows version 23.0 (IBM Corp, USA) was used for the statistical study. The charts were plotted using GraphPad Prism 8.

## 3. RESULTS

### 3.1 The Effect of *P. amarus* Extract on Physical Signs of Toxicity

Weekly observation for overt toxicity symptoms in the rats from the cage sides shows that the animals did not exhibit any sickly signs as presented in Table 1.

### 3.2 The Effect of *P. amarus* Extract on Body Weight Indicators

The result of the *P. Amarus* extract on changes in body weight is depicted in Fig. 1. The oral


administration of the extract caused increase in body weight but with no significant differences ( $p > 0.05$ ) in the body with respect to the control and the respective working doses. Also non-significant changes ( $p > 0.0$ ) was observed with respect to liver weight, relative liver weight, kidney weight, and relative kidney weight, heart weight and relative heart weight.

### 3.3 The Subacute Effect of *P. amarus* Extract on *In-vivo* Antioxidant Enzymes of Male Wistar Rats

The effect of *P. amarus* extract on *in-vivo* antioxidant enzymes of rats in unit/mg tissue is presented in Fig. 2. Findings from the result indicated a significant differences ( $p < 0.05$ ) between the control group and the respective working doses of 1500 and 2000 mg/kg body weight with regards to catalase, superoxide dismutase. Malondialdehyde was also significantly  $p < 0.05$  reduced.

### 3.4 Sub-acute Effects of *P. amarus* Extract on Renal Function Indices of Rats


The effect of *P. amarus* extract on the renal function of rats in mg/dL after 14 days' treatment regimen is presented in Fig. 3. The results obtained indicated a non-significant change ( $p > 0.05$ ) on the concentrations of creatinine, blood urea nitrogen and uric acid.




**Fig 1. Sub-Acute Effects of *P. amarus* Extract on Body Weight Indicators of Male Wistar Rats. Data are mean ± SD of triplicate determinations, values with identical bars with the same superscript letter are not significantly different  $p > 0.05$ . One-way Analysis of Variance (ANOVA)**

**Table 1. Weekly Cage Side Scrutiny for Physical Signs of Toxicity of *P. amarus* Extract on Male Wistar Rats**

S/N	Physical Signs Of Toxicity	DAY 1				DAY 7				DAY 14			
		Control	1000 mg/Kg	1500 mg/Kg	2000 mg/Kg	Control	1000 mg/Kg	1500 mg/Kg	2000 mg/Kg	Control	1000 mg/Kg	1500 mg/Kg	2000 mg/Kg
1	Salivation	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
2	Lacrimation	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
3	Eye dullness	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
4	Eye opacity	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
5	Diarrhea	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
6	Restlessness	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
7	Red stained muzzle	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
8	Lethargy	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
9	Piloerection	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
10	Skin appearance	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal
11	Subcutaneous swelling	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
12	Loss of appetite	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
13	Colour and consistency of faeces	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal	Normal
14	Abdominal distension	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil	Nil
15	Mortality	0	0	0	0	0	0	0	0	0	0	0	0











**Fig. 2. Sub-Acute Effects of *P. amarus* Ethanol Extract on *In-vivo* Antioxidant Status of Male Wistar Rats. Data are mean ± SD of triplicate determinations values with identical bars but with different superscript letters are significantly different  $p < 0.05$ . KEY: CAT-Catalase, SOD- Superoxide dismutase, MDA-Malondialdehyde**



**Fig. 3. Subacute Effects of *P. amarus* Extract on Renal function indices of Rats. Data are mean ± SD of triplicate determinations values with identical bars with the same superscript letter are not significantly different  $p > 0.05$ . One-way Analysis of Variance (ANOVA)**

### 3.5 Histopathology

	Control	1000 mg/kg	1000 mg/kg	1000 mg/kg
<b>LIVER x 40 MAGNIFICATION</b>	 <p>Liver histology appears normal (arrow). Visible centrioles with properly fenestrated sinusoidal space. The hepatocytes appear distinct with well well-differentiated nucleus.</p>	 <p>Liver histology looks very normal (arrow). Visible centrioles with properly fenestrated sinusoidal space. The hepatocytes appear distinct with well well-differentiated nucleus.</p>	 <p>Liver histology appears normal (arrow). Visible centrioles with well fenestrated sinusoidal space. The hepatocytes appear distinct with well well-differentiated nucleus.</p>	 <p>Liver histology appears normal (arrow). Visible centrioles with well fenestrated sinusoidal space. The hepatocytes appear distinct with well-differentiated nucleus</p>
<b>KIDNEY x 40 MAGNIFICATION</b>	 <p>The kidney section shows normal histological features. The section indicated a detailed cortical parenchyma and the renal corpuscles appeared as dense rounded structures (arrow)</p>	 <p>The kidney section shows normal histological features. The section indicated a detailed cortical parenchyma and the renal corpuscles appeared as dense rounded structures (arrow)</p>	 <p>The kidney section shows normal histological features. The section indicated a detailed cortical parenchyma and the renal corpuscles appeared as dense rounded structures (arrow)</p>	 <p>The kidney section shows normal histological features. The section indicated a detailed cortical parenchyma and the renal corpuscles appeared as dense rounded structures (arrow)</p>

### 4. DISCUSSION

Since primeval eras, nutraceutical products, like medicinal plants, have been the foundation for the treatment and management of diver's ailments. This study tries to reveal the effects of subacute doses of *P. amarus*, one of several natural products specifically distributed in the South geological zone of Nigeria [7]

"Alterations in body weight is a salient indicator of the overall health status and well-being of animals" [21]. The results obtained showed that all the animals in the respective experimental

groups underwent an increase in body weight after the treatment regimen. This increase in body weight of the experimental groups administered with graded doses (1000, 1500 and 2000 mg/kg) of *P. amarus* was not significantly different ( $p > 0.05$ ) from the non-treated (control) group, an indication that *P. amarus* at the dose levels investigated did not alter normal metabolic activities of the experimental animals. Flavonoids and lignin in *P. amarus* have been associated with the improvement of glucose, and lipid metabolism, as well as insulin sensitivity [22,23]. The abundant presence of lignans (phyllanthin and hypophyllanthin), flavonoids (quercetin and

rutin), tannin (ellagitannins and gallotannins), phenolic compounds (gallic acid and geraniin) and some alkaloids have been ascribed with potent anti-inflammatory and antioxidant properties [24]. The anti-inflammatory and antioxidant properties of these phytochemicals have been correlated with the maintenance of optimal body weight [25].

“The kidney is actively involved in the routine metabolism of the cell utilizing its role in the excretion of waste products and toxins like urea, uric acid and creatinine, metabolic control of extracellular fluid volume, serum osmolality and electrolyte balance, combined with the production of hormones like erythropoietin and 1,25 dihydroxy vitamin D and renin” [26,27]. “Assessment of renal function markers is vital to the diagnosis, management and treatment of patients with kidney disorders. Pathological conditions affecting renal function may sometimes arise from the ingestion of certain synthetic drugs used in the management and treatment of other underlying health conditions” [19,28]. “Urea is the principal nitrogenous waste produced during protein metabolism and whose level in the blood is reliant upon the correlation between its production and excretion, increased levels beyond the reference range may suggest kidney disease, shock, dehydration, diabetes, acute myocardial infarction while a decreased value lower than normal may portend liver failure, impaired absorption and overhydrating” [29]. Creatinine is the breakdown product of creatine phosphate, primarily from muscle metabolic activities and then excreted by glomerular filtration during normal renal function. Higher values of creatinine above 1.5 mg/dL are pointers of impairment in liver function. Uric acid is a breakdown metabolite from purine metabolism, abnormally high levels of uric acid are associated with a condition called gout. The non-significant changes ( $p > 0.05$ ) in values of urea, creatinine and uric acid by the various doses under investigation when compared to the control, is a positive clue that *P. amarus* ethanol extract had no adverse effect on the renal function indices of the rats. This positive stimulatory effect on the kidney is attributed to the high levels of antioxidant peptides in *P. amarus* [30,31] and other phytonutrients reported in literatures [32,33].

“Oxidative stress arises when the equilibrium between reactive oxygen species (ROS) formation and detoxification promotes an increase in ROS levels leading to agitated

cellular function. ROS causes injury to cellular components leading to lipid peroxidation, nucleic acid, and protein modifications. The formation of lipid peroxidation and the subsequent alteration of nucleic acid and protein are primary etiological factors in the initiation and progression of various metabolic and neurodegenerative diseases” [21,34]. “Oxidative stress is correlated with disturbed redox control mechanisms and cellular signalling pathways, leading to the formation of various forms of cancer and oncogenic initiation and propagation. Antioxidants in cells include catalases, superoxide dismutases (SOD), and glutathione peroxidases (GPX), their induction is usually in response to specific toxicants and pollutants that can induce oxidative stress. Superoxide dismutase (SOD) is essential in protecting cells from oxidative damage by catalysing the dismutation of superoxide radicals into oxygen and hydrogen peroxide, which other antioxidant enzymes like catalase and glutathione peroxidase can further detoxify” [35]. This act is key in preserving redox homeostasis and alleviating oxidative stress, which is implicated in various diseases and ageing progressions. Catalase is an important enzyme that plays a crucial function in defending cells from oxidative damage by disintegrating hydrogen peroxide, a toxic derivative of numerous metabolic activities, into water and oxygen. This reaction is vital in inhibiting oxidative damage to cellular components such as DNA, proteins, and lipids [36]. The significant increase ( $p < 0.05$ ) in the activity of the SOD and catalase by *P. amarus* in this study is key to its optimization of cardiovascular health [37], reproductive health [38], neuroprotection [39], anti-ageing and skin health [40] reported in literature.

Malondialdehyde is a reactive organic compound and a byproduct of lipid peroxidation, it occurs when reactive oxygen species (ROS) attack polyunsaturated fatty acids in cell membranes. It is extensively employed as a biomarker to evaluate the level of oxidative stress in biological systems. Elevated levels of MDA imply increased lipid peroxidation and oxidative damage, which are correlated to numerous disease conditions and ageing [41]. The significantly decreased ( $p < 0.05$ ) MDA of the rats treated with 1500 and 2000 mg/kg of *P. amarus* is a positive indication that the extract has a special ability to mitigate lipid peroxidation.

Histological investigation of the overall architecture of the liver of all groups studied



shows that the integrity of the hepatic lobules was well maintained, with clear demarcation between lobules, an indication of normal liver function and the absence of pathological changes such as steatosis (fatty liver), inflammation, fibrosis, or cirrhosis. The overall architecture of the kidneys of the rats in the respective groups also revealed a very distinct cortex and medulla well-delineated with an appropriate cortical-medullary ratio with the kidney capsule being thin and intact without any signs of thickening or inflammation. Kidney histology reflects the absence of pathological changes such as glomerulosclerosis, tubular atrophy, interstitial fibrosis, and vascular changes, which are indications of several kidney diseases. *P. amarus* demonstrates significant histopathological advantages, chiefly through its antioxidant, anti-inflammatory, hepatoprotective, and nephroprotective activities. These properties make it a valuable therapeutic agent in managing various pathological conditions [42].

## 5. CONCLUSION

This study investigated the toxicological profile of the crude ethanol extract of *P. amarus*. The findings submit that the daily single administration at doses of 1000, 1500 and 2000 mg/kg body weight for two weeks is nontoxic to the albino rats, as depicted by no visible and noticeable overt signs of toxicity, non-significant ( $p > 0.05$ ) adverse change in body and relative organ weight, renal function and organ histopathology of the rats. *P. amarus* also significantly boasts *in-vivo* antioxidant status while significantly reducing the level of malondialdehyde.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

Etiquettes for the use of these animals were endorsed by the Directorate of Research and Quality Assurance, Federal University Otuoke, Bayelsa State vied an approval DRQA/FUO/0100/13/03/24 and the its guidelines were meticulously adhered to.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) now declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image

generators have been used during the writing or editing of this

## COMPETING INTERESTS

The authors have declared that no competing interests exist.

## REFERENCES

1. Kpomah ED, Monday DA, Kpomah B. GCMS analysis of leaves and seeds of *Piper guineense* Schumach & Thoon. African Scientist. 2019;20(3):127-138.
2. Shaito A, Thuan DTB, Phu HT, Nguyen THD, Hasan H, Halabi S, Abdelhady S, Nasrallah GK, Eid AH and Pintus G. Herbal Medicine for Cardiovascular Diseases: Efficacy, Mechanisms, and Safety. Frontiers in Pharmacology. 2020; 11:422. DOI: 10.3389/fphar.2020.00422
3. Kpomah ED, Onyeike EN Kpomah B. Evaluation of some elemental, bioactive compounds and proximate composition of three commonly used herbal plants in the Niger Delta Region of Nigeria. Chemistry Research Journal. 2018;3(2):12-21.
4. Niyomchan A, Chatgat W, Chatawatee B, Keereekoch T, Issuriya A, Jaisamut P, Chusri S, Kunworarath N. Safety Evaluation of the Polyherbal Formulation NawaTab: Acute and Subacute Oral Toxicity Studies in Rats. Evidence-Based Complement Alternative Medicine. 2023:9413458. DOI: 10.1155/2023/9413458. PMID: 37528898; PMCID: PMC10390268.
5. Kpomah ED, Kpomah B, Arhoghro EM. Histomorphological and Biochemical Changes Induced in Male Wistar Rats by Chronic Oral Doses of *Piper guineense* Schumach. & Thonn. Nigerian Journal of Pharmaceutical and Applied Science Research. 2018;7(1):44-51
6. Bose MGA, Banerjee A, Chattopadhyay S. An insight into the potent medicinal plant *Phyllanthus amarus* Schum. and Thonn. Nucleus. 2022; 65,437–472. Available:https://doi.org/10.1007/s13237-022-00409-z
7. Kpomah ED, Ogbogbo J, Kpomah B. Subacute toxicity studies of *Phyllanthus amarus* on haematological parameters and some plasma enzyme activities in mice. International Journal of Basic Science and Technology. 2017;3(1):53-58.

8. Tahseen M, Mishra G. Ethnobotany and diuretic activity of some selected Indian medicinal plants. *The Pharma Innovation*. 2013; 2:112.
9. Kpomah B, Egboh SHO, Agbaire PO, Kpomah, ED. Metal complexes of acetone thiosemicarbazone: synthesis, spectral characterization and pharmacological studies. *Journal of Pharmacological and Applied Chemistry*. 2016;2(2):45-51.
10. Kpomah B, Kpomah, ED, Enemose EA. Activity of some complexes containing 1, 10 Phenanthroline and Thiosemicarbazone derivatives on *Plasmodium Berghei* Infected Strains of Mice, *Nigerian Journal of Applied Sciences*. 2018; 36:13-23.
11. Kpomah ED, Arhoghro EM. Investigation into the intake of a popular polyherbal drug (Jalin Herbal Mannex Liquid) on selected biochemical indices of male wistar rats. *African Journal of Biochemistry Research*. 2022;16(4):55-62.
12. Cohen GD, Dembiec, Marcus J. Measurement of catalase activity in tissue extracts. *Analytical Biochemistry*.1970;34:30-38.
13. Misra HP, Fridovich I. The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *Journal of Biological Chemistry*. 1972;247(10):3170-3175.
14. Hunter FE, Gebicki JM, Hoffsten PE, Weinstein J Scott A. Swelling and lysis of rats' liver mitochondria induced by ferrous ions. *Journal of Biological Chemistry*.1963;23: 828-835.
15. Gutteridge JM, Wilkins S. Copper dependant hydroxyl radical damage to ascorbic acid; formation of thiobarbituric acid reactive products. *FEBS Letters*.1982;137:327-330.
16. Tobacco A, Meiattini F, Moda E, Tarli P. Simplified enzymic/colourimetric serum urea nitrogen determination. *Clinical Chemistry*.1979;25:336-337.
17. Bartels H, Bohmer M, Heuerli, C. Micro-determination of creatinine. *Clinica Chimica Acta*. 1971; 32:81-85.
18. Duncan P, Bayse D, Burnett R, Carey N, Carter R, Fellows WD, Garber C, Kessler G, McComb R, Miller W, Nast P, Ryan W, Schaffer R, Tejada B, Vanderlinde R and Widdowson G. A candidate reference method for uric acid in serum. II. Inter-laboratory testing. *Clinical Chemistry*. 1982; 28:291.
19. Kpomah B, Kpomah ED. Toxicological Assessment of Acetone Thiosemicarbazone Metal Complexes on Body Weight, Biochemical Parameters and Liver Histology of Wistar rats. *Journal of Pharmaceutical and Applied Chemistry*. 2017;3(3):215-224
20. Kpomah B, Obaleye JA, Enemose EA, Kpomah, ED. Cu(II) and Cd(II) Complexes containing 1,10-phenanthroline and methylketone thiosemicarbazone: Synthesis, characterisation and biological activity. *Ife Journal of Science*. 2019;21(3):157-167.
21. Kpomah ED, Arhoghro EM. Positive stimulatory potentials of Coconut (*Cocos nucifera L.*) juice extract on *in-vivo* antioxidants, renal function and lipid profile of male wistar rats. *European Journal of Medicinal Plants*. 2023;34(3):45-54
22. Kpomah B, Kpomah ED, Idu TE, Ugbune U. Mixed ligand complexes containing 2, 2'-Bipyridine with Acetaldehyde Thiosemicarbazone: Synthesis, Characterization and Antifungal Activity. *International Journal of Basic Science and Technology*. 2020; 6(1):26-34
23. Patel P, Shah M. Anti-obesity effect of *Phyllanthus amarus* on high-fat diet-induced obese rats. *Journal of Traditional and Complementary Medicine*. 2017;7(3):386-392.
24. Kpomah ED, Arhoghro EM. Effects of doses of *Bryophyllum pinnatum* and glibenclamide on serum glucose and lipid profile in alloxan-diabetic rats. *Indian Journal of Drugs and Disease*. 2012;1(5):124-128.
25. Patel JR, Tripathi P, Sharma V, Chauhan NS, Dixit VK. *Phyllanthus amarus*: Ethnomedicinal uses, phytochemistry and pharmacology: A review. *Journal of Ethnopharmacology*. 2011;138(2):286-313.
26. Harikrishnan H, Jantan I, Haque MA, Kumolosasi E. Anti-inflammatory effects of *Phyllanthus amarus* Schum. & Thonn. through inhibition of NF-κB, MAPK, and PI3K-Akt signaling pathways in LPS-induced human macrophages. *BMC Complementary and Alternative Medicine*. 2018;18(1):224.
27. Teichman K, Kotfis K, Sienko J. Biomarkers and mechanisms of oxidative stress 20 years of research with an emphasis on kidney damage and renal transplantation. *International Journal of Molecular Sciences*. 2021;22(15):8010.

- Available: <http://dx.doi.org/10.3390/ijms22158010>
28. Kpomah ED, Kpomah B. Microstructural Tissue Assessment, Sex Hormones and Biochemical Investigations following Acute Administration of *Piper guineense* Schumach & Thonn. on female *Rattus norvegicus*. IOSR Journal of Applied Chemistry. 2018;11(5):9-17.
  29. Kpomah B, Kpomah ED, Ugbune U. Transition metal complexes with N, S' And N, N' bidentate mixed ligand: Synthesis, characterization and activity. Applied Science Reports. 2018;22(2):38-44.
  30. Azab AE, Albasha MO, Elsayed ASI. Prevention of nephropathy by some natural sources of antioxidants. Yangtze Medicine. 2017; 1:235-266.
  31. Niazi RK, Malik MN. *Phyllanthus amarus*: A promising herb for nephroprotection. Journal of Nephrology & Therapeutics. 2022;12(3):112-118.
  32. Cesar, APC, Lopes, FES, Azevedo FFN. Antioxidant peptides from plants. Phytochemical review. 2024; 23:95-100. Available:<https://doi.org/10.1007/s11101-023-09875-y>
  33. Ali H, Khan T. Potential of *Phyllanthus amarus* in treating chronic kidney disease. International Journal of Herbal Medicine. 2023;11(4):202-210.
  34. Srinivasan K, Varghese R. Protective effects of *Phyllanthus amarus* in renal disorders. Phytomedicine Research Journal. 2024;15(1):45-52.
  35. Evuen UF Kpomah ED. Comparative impact of solvent extracts of *Spondia mombi* leaves on in-vitro antioxidant and acetylcholinesterase inhibitory activities. Asian Journal of Research in Biochemistry. 2023;12(3):20-29.
  36. Omeje KO, Ezema BO, Onaebi CN. HPLC fingerprint of flavonoids, enzyme inhibition and antioxidant activity of *Newbouldia laevis* stem-bark: an in vitro and in silico study. Future Journal of Pharmaceutical Sciences. 2023; 9:36. Available: <https://doi.org/10.1186/s43094-023-00486-0>
  37. Glorieux C, Calderon PB. Catalase, a remarkable enzyme: Targeting the oldest antioxidant enzyme to find a new cancer treatment approach. Biological Chemistry. 2017;398(10):1043-1058.
  38. Remigante A, Cordaro M, Morabito R. Redox Homeostasis and Antioxidant Strategies in the Pathophysiology. Antioxidants. 2024;13: 281. Available:<https://doi.org/10.3390/antiox13030281>
  39. Shuji T, Norihiro S, Shiro K, Yoshiaki Y, Yasuhiko N, Hiroshi K, Rescue of the Corpus Luteum and an Increase in Luteal Superoxide Dismutase Expression Induced by Placental Luteotropins in the Rat: Action of Testosterone Without Conversion to Estrogen, Biology of Reproduction. 2000; 62(2):398–403, Available:<https://doi.org/10.1095/biolreprod.62.2.398>
  40. Kumar S, Theis T, Tschang M, Nagaraj V, Berthiaume F. Reactive Oxygen Species and Pressure Ulcer Formation after Traumatic Injury to Spinal Cord and Brain. Antioxidants. 2021; 10(7):1013. Available:<https://doi.org/10.3390/antiox10071013>
  41. Flohe L. Looking Back at the Early Stages of Redox Biology. Antioxidants. 2020; 9(12):1254. Available:<https://doi.org/10.3390/antiox9121254>
  42. Ayala A, Muñoz MF, Arguelles, S. Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. Oxidative Medicine and Cellular Longevity. 2014; 360438.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/119874>