



The Inhibition Potentials of Different Honey against *Staphylococcus aureus*, *Escherichia coli* and *Bacillus species* Isolated from Clinical Source

Akani, Nedie Patience¹, Amadi Wemedo, Samuel¹
and Njoku, Onyedikachi Egbuchulem^{1*}

¹Microbiology Department, Rivers State University, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author ANP designed the study and performed the statistical analysis. Author NOE wrote the first draft of the manuscript and managed the literature searches. Author AWS managed the analyses of the study. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPR/2020/v5i230131

Editor(s):

(1) Dr. Khadiga Ahmed Ismail Eltris, Ain Shams University, Egypt.

Reviewers:

(1) Karima Mogahed Fahim, Cairo University, Egypt.

(2) Melania Florina Munteanu, "Vasile Goldiş" Western University of Arad, Romania.

(3) Srikanth Naramala, Adventist Medical Centre, USA.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/61426>

Original Research Article

Received 25 July 2020

Accepted 30 September 2020

Published 02 November 2020

ABSTRACT

As a result of the increased prevalence of antibiotic resistance among different bacteria, different plants and other natural products have been studied and found to be highly effective against pathogenic bacteria. Honey, over the years has been used as an antibacterial agent to treat certain infections caused by bacteria and is believed to be effective especially in rural areas. This study was thus aimed at comparing the effect of different honey samples against some pathogenic bacteria (*Escherichia coli*, *Staphylococcus aureus* and *Bacillus cereus*) isolated from clinical source. This study was carried out in the microbiology laboratory, department of microbiology Rivers State University Nigeria from January 2018 to August 2019. The antibacterial sensitivity test was carried out using agar well diffusion method while the Minimum inhibitory concentration and Minimum bactericidal concentration were determined using broth tube micro dilution technique in two fold dilution. The inhibition efficiency of the honey samples on the test organisms increased with increasing concentration from 20 to 100% as 100% concentration had the highest zone of

*Corresponding author: Email: finest.blessingd@gmail.com;

inhibition. *Staphylococcus aureus* (6.33 mm – 26.33 mm) was the most sensitive to the honey samples while *Bacillus cereus* (0.00 – 19.67 mm) was less sensitive. At concentrations of 20 – 80%, raw and Rowse honey were more effective on *E. coli* compared to Princenic Global honey, while at 100%, Princenic Global honey was more effective on *Staphylococcus aureus*. Raw and Rowse honey were more effective at 20 -60% concentrations followed by Princenic Global honey; whereas at 80 -100% concentrations, Raw and Princenic Global honey were more effective. *Bacillus cereus* was resistant to the honey samples at 20 – 60% but sensitive at 80 – 100% concentrations to Rowse, Raw and Princenic Global honey. The inhibition efficiency of the honey samples on the growth of the tested organisms was found to be dependent on the concentration and type of honey used, as well as they type of organism tested. The result of the minimum inhibitory and minimum bactericidal concentration showed that *Staphylococcus aureus* was inhibited most at a lower concentration of 25% compared to other bacteria isolates. All honey samples tested did not show any bactericidal effect but was bacteriostatic to some of the tested organisms. Pharmacological standardization and clinical evaluation on the effect of honey is essential before honey can be used as a preventive and curative measure to common diseases related to the tested bacterial species.

Keywords: Honey; antibiotic resistance; antimicrobial agents; minimum inhibitory concentration.

1. INTRODUCTION

European Union Council Directive defined honey as the natural sweet substance produced by *Apis mellifera* bees from the nectar of plants or from the secretions of living parts of plants or excretions of plant sucking insects on the living parts of plants which bees collect, transform by combining with specific substances of their own, deposit, dehydrate, store and leave in honey combs to ripen and mature [1]. Bogdanov in their research stated that honey is the only food sweetener that can be used industrially without processing. Honey can be classified according to its origin (such as nectar or honey dews), mode of production and preservation [2].

Honey as a concentrated aqueous solution composed of a mixture of glucose and fructose but also contains at least 22 other complex carbohydrates, various amino and organic acids, proteins, enzymes, phenol antioxidants, vitamins, minerals, pigments, waxes and pollen grains [3]. In nature honey is very viscous and acidic ranging between 3.2 and 4.5 in pH. Over the years around the world, honey has been effectively used as medicine most especially as traditional remedy in so many countries. Most of the ancient countries that have used honey as a traditional remedy include Egyptians, Assyrians, Chinese, Greeks and Romans. This countries employed honey for wounds and diseases of the gut [4]. Over the years, many researchers have reported the antibacterial activity of honey and found that natural unheated honey has broad-spectrum antibacterial activity when tested against some pathogenic and oral bacteria [5]. In

some developed countries, honey is used for the treatment of ulcers, bed sores and other skin infections resulting from burns and wounds [6].

Lusby in their research indicated that the healing properties of honey can be attributed to the fact that it offers antibacterial activity, keeps the wound environment moist which promotes fast healing and has a glueyness which helps to provide a defensive barrier to prevent infection [7]. Many researchers have shown that the properties present in honey that is responsible for the antibacterial activity is the phytochemical properties such as high content of reducing sugar, high viscosity, high osmotic pressure, low pH, low water activity, low protein content and presence of hydrogen peroxide [8]. Again Alnimat stated that the main antibacterial agent in honey is hydrogen peroxide, which is produced by glucose-oxidase action [9]. The level of hydrogen peroxide in honey is determined also by the presence of catalase, which originates from the pollen of plants. Light, temperature and oxygen affect the amount of hydrogen peroxide which shows a discrepancy according to the processing and storage conditions of the honey. Research has revealed a positive correlation between the endogenous hydrogen peroxide concentration and the inhibitory activity of bacterial growth by honey [10]. Indeed honey with a high concentration of hydrogen peroxide has a higher antibacterial activity.

Libonatti and his group reported that the antibacterial activity of honey is due entirely to the non-peroxide components such as acidity,

osmolarity, flavonoids, phenolic compounds and lysozyme [11]. Different studies have claimed that honey contains bioactive components such as lysozyme, a well-known antibacterial agent [12].

Abd-El Aal when comparing the inhibitory activity of honey and some commonly used antimicrobial agents on some gram-negative bacteria (*Pseudomonas aeruginosa*, *Enterobacter* spp, and *Klebsiella* spp.) showed that honey had a pronounced inhibitory activity of 85.7%. A 100% inhibition was observed in the case of gram positive methicillin resistant *Staphylococcus aureus* in comparison to the use of antibiotics alone [13].

Kwakman and Zaat reported that the sugar content of honey is sufficient to retain antibacterial activity when diluted to approximately 20-40%. Based on extensive research on the medicinal uses of honey, antimicrobial action of honey on *Staphylococcus aureus*, *Escherichia coli* and *Bacillus cereus*, isolated from open wound were investigated [14].

2. MATERIALS AND METHODS

2.1 Materials

2.1.1 Collection and preparation of samples

A total of three honey samples (Princenic Global, Rowse and Raw) were used in this study. Princenic Global (PG) and Rowse honey were bought from supermarket in Port- Harcourt metropolis and Raw honey was bought from local bee keepers in Etche local Government Rivers state, Nigeria. The samples were stored in sterile bottles at temperature of 20 – 21°C in a dark place before analyses. These honey samples were selected because they are widely sold in the supermarkets around the area and they are regularly consumed by people around the area.

2.1.2 Collection and confirmation of bacteria isolates

Bacterial isolates used in this study were wound associated bacteria including *Staphylococcus aureus*, *Escherichia coli* and *Bacillus cereus*. The isolates were obtained from Optimal Diagnostic Center, Mgbuoba, Port-Harcourt. The isolates were collected in sterile Bijou bottles containing nutrient broth and immediately incubated at a temperature of 37°C for 24 hours. The isolates were identified microscopically and biochemically

using tests such as Grams Stain, catalase, simmon Citrate utilization, indole, motility, methyl Red-Voges proskauer, oxidase, sugar fermentation, starch hydrolysis, coagulated, hemolysis and spore stain.

2.2 Methods

2.2.1 Antibiotics sensitivity test

This test was performed using disc diffusion method [15]. The test organisms were first standardized. Five colonies from fresh 24 hours culture were aseptically transferred to a 4 ml sterile normal saline, the suspension was compared with 0.5 McFarland prepared by adding 0.05 ml of 1% Barium Chloride (BaCl) to 9.95 ml of 1% sulfuric acid. The standardized isolates were then streaked on the surface of a large Mueller-Hinton agar plate and allowed to dry for 5 min. Eight (8) commercially-prepared, fixed concentrations, Abtek paper antibiotic disks were placed on the inoculated agar surface using sterile forceps. The antibiotics used include; Ceftazidime (30 µg), Cefuroxime (30 µg), Gentamicin (10 µg), Ceftriaxone (30 µg), Erythromycin (5 µg), Cloxacillin (5 µg), Ofloxacin (5 µg), Augmentin (30 µg), Cefixime (5 µg), Nitrotrantion (300 µg), Ciprofloxacin (5 µg). Plates were incubated at 37°C for 24 hours. After incubation the zones of growth inhibition around each of the antibiotic disks was measured to the nearest millimeter. The zone diameters of each drug are interpreted using the criteria published by the Clinical and Laboratory Standards Institute [15].

2.2.2 Antibacterial sensitivity test of honey

The antibacterial activity of honey samples was tested *in vitro* against three pathogenic bacteria (*Staphylococcus aureus*, *Escherichia coli* and *Bacillus cereus*) using agar well diffusion method [16,17]. The honey samples were prepared by diluting each in sterilized distilled water at different dilutions (concentration), 20%, 40%, 60%, 80% and also net honey (100%). Concentrations were achieved by adding 80 ml of distilled water to 20 ml of honey sample (20%), 60 ml of distilled water to 40 ml of honey, 40 ml of distilled water to 60 ml of honey, 20 ml of distilled water to 80 ml of honey, and lastly 100% was pure undiluted honey.

Mueller-Hinton agar plates were prepared and each plate was properly inoculated with each test organism using a sterile swab stick dipped into the inoculum suspension and spread over the

surface of the medium. Wells were made using a sterile cork borer and each well was filled with different concentrations of the honey. A distance was maintained from the edges of the plates to prevent overlapping of the inhibition zones. The plates were incubated for 24 hrs at 37°C. After incubation the plates were examined and the diameter of the inhibition zones was measured in triplicate for each isolate.

2.2.3 Minimum Inhibitory Concentration (MIC)

The minimum inhibitory concentration of the honeys was determined using broth tube micro dilution method [18]. The purpose of this test was to determine the minimum concentration of honey that can inhibit growth of the test organisms. Eight (7) sterile test tubes were placed in a rack and labeled 1 to 5. Honey control tube (HC) and growth control tube (GC) were used as quality controls. One ml (1 ml) of freshly prepared nutrient broth was added to each tube, sterilized and cooled. Then 1 ml of undiluted honey solution (100%) was added to test tube number 1 and honey control with a sterile micropipette and tips. Then serial twofold dilution was performed by transferring 1 ml undiluted honey into the second tube with separate sterile micropipette. After a thorough mixing, 1 ml of the honey sample was transferred with another sterile micropipette from tube 2 and tube 3. This procedure continued until six tubes with a dilution of 1:125 was reached and finally 1 ml was taken and discarded from tube 5. The growth control tube received no honey was served as a growth control while the HC tube received no bacterial inocula served as honey control.

Except the honey control tube, each tube was inoculated with 1 ml of the culture of respective prepared organism. The procedure was repeated for all the organisms tested to each of the honeys. Tubes were then incubated at 37°C for 24 hours and observed by visual inspections for the presence and absence of growth (turbidity).

2.2.4 Minimum Bactericidal Concentration (MBC)

The test was done to determine the minimum concentration of honey that can kill the test isolates. To determine the MBC, incubated tubes showing no visible sign of growth/turbidity in MIC, were sub-cultured onto sterile nutrient agar plates by streak plate method and incubated at 37°C for 24 hours aerobically. The least

concentration of honey that did not show growth of test organisms was considered as the MBC [18]. Then inoculated plates were scored as bactericidal if no growth; bacteriostatic if there is light to moderate growth and no antibacterial activity if there is heavy growth [18].

2.3 Statistical Analysis

Results obtained were expressed as mean \pm standard deviations and differences between means were analyzed statistically using analysis of variance (ANOVA) on the SPSS version 22.0; differences were considered significant when $p < 0.05$ and where differences occurred, Tukey method was used to separate the means.

3. RESULTS AND DISCUSSION

Table 1 showed the antibiotics sensitivity of the bacteria isolates. Ofloxacin showed the highest zone of clearing on *Staphylococcus* while Cloxacillin and ceftazidime did not have any effect on *Staphylococcus aureus*, this means that the organism was resistant to Cloxacillin and ceftazidime. Again for *Bacillus cereus*, Ofloxacin also showed the highest zone of inhibition of 25 mm while Cloxacillin and ceftazidime showed no zone of clearing meaning the organism was resistant to these antibiotics. *Escherichia coli* were resistant to Augmentin and Cefuroxime while Ciprofloxacin was more effective with inhibition zone of 24.7 mm.

Fig. 1 showed the effects of different concentrations of honey samples on the growth of *Escherichia coli*. From this Figure, PG honey at concentrations of 20, 40, 60, 80 and 100% gave 0.00, 3.67, 12.00, 20.33 and 31.00 mm zones of inhibition, respectively. Also raw honey at the same concentrations gave 0.00, 6.67, 17.67, 22.67 and 29.33 mm zones of inhibition, respectively while Rowse honey at similar concentration on *Escherichia coli* gave 1.33, 8.33, 15.33, 21.67 and 29.67 mm zones of inhibition, respectively.

Fig. 2 showed the effect of different concentrations of honey samples on the growth of *S. aureus*. It showed that PG honey gave zones of inhibition of 6.33, 10.33, 15.67, 21.00 and 28.67 mm at concentrations of 20, 40, 60, 80 and 100%, respectively while on Raw honey at similar concentration gave zones of inhibition of 8.00, 13.00, 18.67, 23.67 and 29.67 mm, respectively. Furthermore, the effect of Rowse

honey on the growth of *S. aureus* at concentrations of 20, 40, 60, 80 and 100% gave 7.67, 10.00, 15.33, 18.33 and 26.33 mm zones of inhibition, respectively.

Table 1. Antibiotics sensitivity pattern exhibited by test bacterial isolates

Isolates	Ofloxacin(5µg)	Augmentin(30 µg)	Nitrotrantion(30 µg)	Ciprofloxacin(5 µg)	Ceftazidime(30 µg)	Cefuroxime(30 µg)	Gentamycin(10 µg)	Cefixime(5 µg)	Ceftaxone(30 µg)	Erythromycin(5 µg)	Cloxacillin(5 µg)
<i>Escherichia coli</i>	S	R	S	S	I	R	S	S	-	-	-
<i>Staphylococcus Sp</i>	S	S	-	-	R	S	S	-	S	S	R
<i>Bacillus Sp</i>	S	I	-	-	R	I	S	-	S	S	R

Keys: R = resistance (≤ 13)
 S = sensitive (≥ 17)
 I = intermediate ($\geq 14-16$)




Fig. 1. Effects of different concentrations of honey samples on the sensitivity pattern of *Escherichia coli* as shown by the diameter of the zones of inhibition




Fig. 2. Effects of different concentrations of honey samples on the sensitivity pattern of *Staphylococcus aureus* as shown by the diameter of the zones of inhibition

Fig. 3 showed that only 80 and 100% concentrations of the honey samples were effective on *B. cereus*. PG honey was effective on *B. cereus* at a concentration of 100% with zone of inhibition of 1.33 mm, while raw honey inhibited its growth at 80 and 100% concentrations with zones of inhibition of 2.33 and 7.00 mm, respectively. Also, Rowse honey inhibited at 80 and 100% concentrations with inhibition zones of 9.67 and 14.67 mm, respectively.

Table 2 showed that all the three honey types inhibited the growth of the bacterial isolates at a minimum concentration of 50%. This indicated that the bacterial isolates were sensitive to the honey samples at 50% concentration while resistant to honey samples at the rest concentrations.

Table 3 showed that all the honey samples were not bactericidal to the bacterial isolates except PG honey which was bacteriostatic to *Staphylococcus aureus* at 50%.




Fig. 3. Effects of different concentrations of honey samples on the sensitivity pattern of *Bacillus cereus* as shown by the diameter of the zones of inhibition

Table 2. The minimum inhibitory concentration of honey samples at different concentrations against bacterial isolates

Honey samples	Concentrations of honey (%)														
	<i>E. coli</i>					<i>S. aureus</i>					<i>Bacillus</i>				
	50	25	12.5	6.5	3.1	50	25	12.5	6.5	3.1	50	25	12.5	6.5	3.1
PRINCENIC GLOBAL	S	R	R	R	R	S	S	R	R	R	S	R	R	R	R
ROWSE	S	R	R	R	R	S	S	R	R	R	S	R	R	R	R
RAW	S	R	R	R	R	S	S	R	R	R	S	R	R	R	R

Keys; S = Sensitive, R = Resistant

Table 3. The minimum bactericidal concentration of honey samples against bacteria isolates at different concentration

Honey samples	<i>E. coli</i>		<i>S. aureus</i>		<i>Bacillus</i>	
	50%	25%	50%	25%	50%	25%
PRINCENIC GLOBAL	++		+	++	++	
ROWSE	++		++	++	++	
RAW	++		++	++	++	

Keys; NS= Non-sterilized; ++= not bactericidal (heavy growth), += bacteriostatic (light growth)

3.1 Discussion

Results of this study showed that Rowse honey inhibited the growth of *Escherichia coli* at the lowest concentration compared to other honey types. At 100% PG honey had higher inhibition zone on *Escherichia coli* with zone in diameter of 31.00 mm, whereas *Staphylococcus aureus* at 20% had inhibition zone of 6.33 mm by PG honey, while *Bacillus* was not inhibited at 20%. Comparison of this honey types showed that *Staphylococcus aureus* was most inhibited by the honey types at lowest concentration followed by *Escherichia coli* and *Bacillus*. Again Rowse honey was most effective at the lowest concentrations against *Escherichia coli* and *Bacillus* while raw honey was most effective against *Staphylococcus aureus* at lowest concentration. The wide-ranging inhibition level of the honey samples is due to the fact that different honey types possess different efficacies against the same type of bacterium and different bacteria [19]. Reports have shown that the ability of honey to inhibit microbial growth is not only due to osmolality, viscosity, presence of hydrogen peroxide and low protein contents but also due to other factors that affect the composition of honey [19]. Such factors depend on a great extent on the bee's source, the location of the flowers and related weather conditions, the storage time and conditions and the method of preservative treatment [20].

The results of this study were in agreement with the study performed by Al-Haj who used Malaysian honey on both methicillin sensitive *Staphylococcus aureus* and methicillin resistant *Staphylococcus aureus*. They concluded that honey completely inhibited the growth of the two bacteria [21]. Also, the reports of this study is in consonance with the study by Taormina, where they investigated the antibacterial activity of honey from six floral sources against *Escherichia coli*, *Salmonella thyphimurium*, *Shigella sonnei*, *Staphylococcus aureus* and *Bacillus cereus* using disc diffusion method. Their results showed that the development of inhibition zones

depended on the concentration of the honey used as well as the test pathogen; their result shows that *B. cereus* was least inhibited while *S. aureus* was most inhibited by the different honey samples [22].

4. CONCLUSION

The results of this study showed an increase in concentrations of honey samples increased their inhibitory effects on the test isolates. Also, among the three studied pathogenic bacteria, *E. coli* was the most inhibited with 29.33, 29.67 and 31.00 mm zones of inhibition by Raw, Rowse and PG honey samples, respectively while *B. cereus* was the least inhibited with 1.33, 7.00 and 14.67 mm zones of inhibition by PG, raw and Rowse honey samples, respectively. Comparison of the results of the figures showed that PG honey was most effective on *Escherichia coli* with zone a inhibition of 31.00 mm while on *S. aureus* Raw honey was the most effective with a diameter of 29.67 mm. Also, Rowse honey showed higher efficiency on *B. cereus* with inhibition diameter of 14.67 mm. Although, the three honey samples exhibited varied inhibitory effects on the same bacterium and the different bacteria, all three samples were found to have antibacterial effects against the isolates. This further proves that honey is a potent antibacterial agent and could be used in place of synthetic antibiotics if properly standardized especially with the rising occurrence of antibiotic resistance among synthetic drugs.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. European Union Council Directive. European commission committee meeting of 20th December, 2001 relating to honey. Washington D.C. 2002;25:11-14.

2. Bogdanov S, Ruoff K, persano OL. Physico-chemical methods for characterization of unifloral honey: A review. *Apidologie*. 2004;35:4–17. DOI: 2004047
3. Bogdanov S, Haldimann M, Luginbuhl W, Gallmann P. Minerals in honey: Environmental, geographical and botanical aspects. *Journal of Apicultural Research*. 2007;46:269–275. DOI: 10.3896/IBRA.1.46.4.11
4. Bogdanov S, Jurendic T, Sieber R, Gallman P. Honey for nutrition and health. A Review: *Journal of American College of Nutrition*. 2008;27(6):677–689. DOI: 10.1080/07315724.2008.10719745
5. Mavric E, Withmann S, Barth G, Henle T. Identification and quantification of methylglyoxal as the dominant antibacterial constituents of Manuka honeys from New Zealand. *Molecular Nutrition and Food Research*. 2009;52:483–489. DOI: 10.1002/mnfr.200700282
6. Cooper R, Molan P, Harding K. The sensitivity to honey of gram positive cocci of clinical significance isolated from wounds. *Journal of Applied Microbiology*. 2002;93(5):857–863. DOI: 10.1046/j.1365-2672.2002.01761
7. Lusby PE, Coombes AL, Wilkinson JM. Bactericidal activity of different honeys against pathogenic bacteria. *Arch Med Res*. 2005;36:464–467. DOI: 10.1016/j.arcmed.2005.03.038
8. Molan PC, Cooper RA. Honey and sugar as dressing for wounds and ulcers. *Tropical Doctor*. 2000;30:249–251. DOI: 10.1177/004947550003000429
9. Alnaimat S, Wainwright M, Al Bud K. Antibacterial potential of honey from different origins: A comparison with Manuka honey. *Journal of Microbiology, Biotechnology and Food Sciences*. 2012;1:1328–1338. Available: <https://www.researchgate.net/publication/236684213>
10. Bizerra FC, Da Silva PI, Hayashi MAF. Exploring the antibacterial properties of honey and its potential microbiology. *Frontiers Microbiology*. 2012;3(1):398–400. DOI: 10.3389/fmicb.2012.00398
11. Libonatti C, Soledad V, Marina B. Antibacterial activity of honey: A review of honeys around the world. *Journal of Microbiology and Antimicrobials*. 2014;6(3):51-56. DOI: 10.5897/JMA2014.0308
12. Estrada H, Gambia M, Anas M, Chaves C. Evaluation of the antimicrobial action of honey against *Staphylococcus aureus*, *Staphylococcus epidermis*, *Pseudomonas aeruginosa*, *Escherichia coli* and *Aspergillus niger*. *Archivos Latino Americanos de Nutrition*. 2005;55:167–171. PMID: 16335227
13. Abd–El Aal A, El-Hadicy M, El-Mashad N, El-Sebane A. Antimicrobial effect of bee honey in comparison to antibiotics on organisms isolated from infected burns. *Annual Burns Fire Disaster*. 2007;20:83–86. PMC3188067.
14. Kwakman PH, Zaat SA. Antibacterial components of honey. *Life*. 2012;64(1): 48–55. DOI: ORG/10.1002/IUB.578
15. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial disk susceptibility tests. Approved Standard M2-A102009 Wayne, PA Clinical and Laboratory Standards Institute; 2013.
16. Cooper R, Molan P, Harding K. The sensitivity to honey of gram positive cocci of clinical significance isolated from wounds. *Journal of Applied Microbiology*. 2002;93(5):857–863. DOI: 10.1046/j.1365-2672.2002.01761
17. Okeke O, Okeke MU, Ezejiofor CC, Ndubuisi JO. Antimicrobial activity of honeys from Nsukka and Ugwuaji in Enugu State, on selected pathogenic bacteria isolated from wound. *Advances in Analytical Chemistry*. 2018;8(1):6-9. DOI: 10.5923/j.aac.201808
18. Kacaniova M, Vukovic N, Bobkova A, Fikselova M, Rovna K, Hascik, P. Antimicrobial and antiradical activity of Slovakian honeydew honey samples. *Journal of Microbiology Biotechnology and Food Sciences*. 2011;1(3):354–360. Available: <https://www.researchgate.net/publication/313607835>
19. Almasaudi SB, Alea AM, Elsayed M, Yousef AQ, Esan A, Mohamad Q. Antimicrobial effect of different types of honey on *Staphylococcus aureus*. *Saudi Journal of Biological Sciences*. 2017;24: 1255–1261. DOI: 10.1016/j.sjbs.2016.08.007
20. Jing P, Xu W, Yi H, Bai L, Yuan R. An amplified electrochemical optasensor for thrombin detection based on Pseudobioenzyme Fe₃O₄– Au nano composites and electro active hemin/G-

- quadruplex and signal enhances. *Analyst*. 2014;139:1756–1761.
21. Al-Hga NA, Amghalia E, Shamsudin MN, Abudullah R, Sekawi Z. Antimicrobial activity of honey against methicillin resistant *Staphylococcus aureus*. *Research Journal of Biological Sciences*. 2009;4:943–947.
22. Taormina PJ, Niemira BA, Beuchat LR. Inhibitory activity of honey against food borne pathogens and influenced by the presence of hydrogen peroxide and levels of anti-oxidant power. *International Journal of Food Microbiology*. 2001;69: 217–225.
DOI: 10.1016/s0168-1605(01)00505-0

© 2020 Patience et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/61426>