



# Optimizing Goldfish (*Carassius auratus*) Breeding and Larval Nurturing for Materialistic Aquaculture Utilizing Synthetic Hormones

Dhrubajyoti Panja <sup>a++</sup>, Rakhi Das <sup>a##</sup>, Bijay Kali Mahapatra <sup>b†</sup>  
and Deepak Kher <sup>c‡</sup>

<sup>a</sup> Department of Aquaculture, Sanjeev Agrawal Global Educational University, India.

<sup>b</sup> CIFE, Kolkata, India.

<sup>c</sup> School of Agriculture, Sanjeev Agrawal Global Educational University, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/air/2024/v25i41104>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/119519>

Original Research Article

Received: 04/05/2024  
Accepted: 06/07/2024  
Published: 10/07/2024

## ABSTRACT

Gold fish (*Carassius auratus*) are familiar in India as an ornamental freshwater fish with economic importance. This experiment aimed to explore the possibility for mass-scale breeding and larval rearing of goldfish under biofloc setup. Natural brooders were collected, acclimatized and induced

<sup>++</sup> Research Scholar;

<sup>#</sup> Assistant Professor;

<sup>†</sup> Retired Scientist;

<sup>‡</sup> Dean;

\*Corresponding author: E-mail: [rakhi.d@sageuniversity.edu.in](mailto:rakhi.d@sageuniversity.edu.in);

**Cite as:** Panja, Dhrubajyoti, Rakhi Das, Bijay Kali Mahapatra, and Deepak Kher. 2024. "Optimizing Goldfish (*Carassius Auratus*) Breeding and Larval Nurturing for Materialistic Aquaculture Utilizing Synthetic Hormones". *Advances in Research* 25 (4):271-75. <https://doi.org/10.9734/air/2024/v25i41104>.

to breed utilizing synthetic hormones at various dosages. Breeding execution parameters namely spawning success, fertilization rate and hatching performance were analysed. The highest fertilization and hatching rate were gradually noticed at 80.8% and 80.9% by using a dose of 0.4ml/kg body weight for females and 0.2ml/kg for males. Larval rearing was accomplished by applying several live and formulated feeds across various life stages in normal rearing tanks and biofloc tanks also. The growth performance of larvae in biofloc tanks express a satisfactory rate, with a maximum weight gain of 7.1 g and length gain of 5 cm over 120 days compared to 5.1 g and 3.5 cm in normal tanks. Water quality parameters were kept within the optimum range all over the experiment. The functionable study exhibits the viability of captive breeding and mass scale larval nurturing of goldfish employing induced breeding techniques and biofloc technology.

*Keywords: Gold fish; induced breeding; biofloc; production.*

## 1. INTRODUCTION

Today as an ornamental fish species, gold fish are supposed to be the most popular fish which kept in aquariums and outdoor ponds globally. Their vigorous colours, hardy nature and distinctive behaviour have built them a preferred among hobbyists and aqua culturists alike. Beyond their decorative value, goldfish also carry cultural significance in several traditions and have been the subject of particular breeding for centuries, ensuing in diverse captivating varieties [1,2,3].

India is a leading country for production of ornamental fish due to its divergent aquatic ecosystem, suitable climate condition and intensive labour cost [4,5]. With the emerging demand for ornamental fish in the aquarium business, there is a crucial need to develop effective and regenerative production methods for goldfish. Generally healthy and mature gold fish are preferred to meet the demand of commercial market [6,7-9]. Conventional breeding practices frequently rely on natural spawning, which can be uncertain and production inconsistent results. Additionally the survivability and growth rate of goldfish larvae can be initiated by different factors such as water quality, feed composition and rearing conditions [10,11].

This research intended for evaluate the prospective for large-scale breeding and larval rearing of goldfish under captive conditions [12-14]. By applying induced breeding techniques and advanced larval rearing systems, the researchers sought to flourish a reliable and systematic method for producing high quality goldfish on a commercial scale.

## 2. MATERIALS AND METHODS

### 2.1 Collection and Acclimatization of Brooders and Observe Breeding Behaviour

Adult goldfish brooders were collected from local water bodies utilizing normal fish capture techniques. The fish were transported to the research potential and moderately acclimated to captive conditions in specific designed holding tanks. Different physico chemical parameters like temperature, dissolved oxygen, pH, and ammonia levels were sharply monitored and maintained within optimal levels for goldfish. Fishes were kept in separate tank according their sexual variation and monitoring the sexual courtship under the captivity.

### 2.2 Artificial Breeding Setup and Procedures

Brooders were precisely chooses based on their morphological characteristics and reproductive maturity. Both males and females were segregated and nurture in separate spawning tanks equipped with suitable substrates and ecological conditions helpful to breeding.

### 2.3 Hormone Administration Protocols

For getting an expected result from induce spawning, synthetic hormones (HATCHME composition of SGnRH 0.002%, Domperidone 0.998%, Proylene Glycol 99.0%) were applied to the brooders. Distinctive doses synthetic gonadotropin releasing hormone analogue (SGnRHa) was checked to determine the standard dosage for successful breeding. The hormones were applied through intramuscular injections and the brooders were closely monitored for symptoms of spawning.

## 2.4 Larval Rearing Systems

Two well defined larval rearing systems were constructed in this research: normal rearing tanks and biofloc tanks. The normal rearing tanks were furnished with specific aquaculture practices, while the biofloc tanks assimilate the principles of biofloc technology, which entail the promotion of useful microbial communities in the rearing water.

## 2.5 Feed Formulation and Feeding Regimes

Larvae were fed a blend of live and artificial feeds at various stages of their growing period. Live feeds mainly artemia nauplii and rotifers were provided during the primary stages, while transition to artificial diets arise as the larvae grew. The formulated feeds were attentively framed to meet the nutritional demand of goldfish larvae, and feeding regimes were enhanced based on larval growth and development.

## 2.6 Sampling and Data Collection

Uniform sampling was drive to monitor the survivality rate of the larvae, growth rate and water quality parameters. During the experimental period, length and weight measurements were checked out by predetermined intervals and survival rates were calculated by following survivality law which was counting the number of live larvae in each rearing system.

## 2.7 Water Quality Parameters

Several water quality related parameters like dissolved oxygen, pH, ammonia, nitrite and nitrate levels, were checked regularly utilizing standard analytical methods. Adjustments were

assembled as needed to maintain ideal conditions for larval growth and development.

## 3. RESULTS AND DISCUSSION

### 3.1 Observation of Sexual Dimorphism and Breeding Behaviour during Experiment

In this experiment, sexual dimorphism and breeding behaviour of goldfish were recorded through continuous observation. Males express visible breeding coloration with prominent tubercles on the operculum and pectoral fins and female show bit fatter than normal. Especially courtship period, chasing and nudging behaviour was noticed.

### 3.2 Spawning Success, Fertilization and Hatching Rates Across Treatment Groups

The evaluation reported the spawning success, hatching rates, fertilization rates achieved across distinct treatment groups, including different dosages of the synthetic hormones. The highest rate of fertilization and hatching rate were noted at 80.8% and 80.9% by applying the dosage of 0.4ml/kg body weight for females and 0.2ml/kg for males.

### 3.3 Growth Performance (Length and Weight Gain)

Larval growth performance was checked by measuring the length and weight gain over time. The larval growth rate in biofloc tanks show a tremendous growth rate than the normal rearing tanks which was 7.1 g in weight and 5 cm in length at biofloc tanks compared to 5.1g in weight and 3.5 cm in length at normal tank over a 120-day period.



Fig. 1. Difference between male (Left one) and female fish (Right one)

**Table 1. Survivality of Goldfish under Captivity**

Tank size	Tank types	Number of fish	Survivality rate
4''×1.5''×1''	Hatchling Tank	960	81.9%
4''×1.5''×1''	Nursery Tank	787	83.8%
4''×1.5''×1''	Rearing Tank	660	75.7%
4''×2''×1''	Biofloc Tank	120	63.3%

### 3.4 Larval Survival Rates in Different Rearing Systems

The survivality rate of larvae in both normal and biofloc tank were monitored carefully during the research. According to the recorded data the survivality rate was higher in normal tank compared to biofloc tank.

### 3.5 Water Quality Analysis

During the experimental period different physico chemical parameter like temperature, dissolved oxygen, pH, ammonia, nitrite and nitrite levels were maintained both normal and biofloc rearing tanks for getting optimal result from the experiment. Temperature ranges between 20 to 24°C, pH extent from 6-8 and ammonia noted as 0.01 to 0.05 ppm.

## 4. CONCLUSION

The focus of the research was to explore the potential applications of the findings from the experiment in the context of mass scale goldfish production. The recorded data indicate the feasibility of adopting induced breeding techniques and enhanced larval rearing systems for commercial-scale operations of goldfish considering economic viability and sustainability.

### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

We, hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

- Demir O, Sarigoz S. Development of feeding program for early larval stage of

gold fish (*Carassius auratus*). Turkish Journal of Fisheries and Aquatic Sciences. 2016;16(2). Available: [https://doi.org/10.4194/1303-2712-v16\\_2\\_12](https://doi.org/10.4194/1303-2712-v16_2_12)

- Walker HJ, Ingole B, Nayak GN, Wafar M, Wafar S, Yennavar P, Bokuniewicz H, Lambeck K. I. In M. L. Schwartz (Ed.), Encyclopedia of Coastal Science. Springer Netherlands. 2005;542–571. Available: [https://doi.org/10.1007/1-4020-3880-1\\_9](https://doi.org/10.1007/1-4020-3880-1_9)
- Pratiwy, Fitri Meyllianawaty, Dini Agustiani. "The Use of Fermented Feed on the Growth of Tilapia (*Oreochromis niloticus*): A Review". Asian Journal of Fisheries and Aquatic Research. 2023;21(1):43-47. Available: <https://doi.org/10.9734/ajfar/2023/v21i11528>.
- Mahapatra B, Kavungal V, Mandal S. Production of Gold Fish, *Carassius auratus* (Linnaeus) and its Economics in Meghalaya. J. Inland Fish. Soc. India; 2013. Available: <https://www.researchgate.net/publication/261525261>
- Shete AP, Verma AK, Tandel RS, Prakash C, Tiwari VK, Hussain T. Optimization of Water Circulation Period for the Culture of Goldfish with Spinach in Aquaponic System. Journal of Agricultural Science. 2013;5(4):26. Available: <https://doi.org/10.5539/jas.v5n4p26>
- Lorenzoni M, Corboli M, Ghetti L, Pedicillo G, Carosi A. Growth and reproduction of the goldfish *Carassius auratus*: A case study from Italy. In F. Gherardi (Ed.), Biological invaders in inland waters: Profiles, distribution, and threats. 2007;2:259–273. Springer Netherlands. Available: [https://doi.org/10.1007/978-1-4020-6029-8\\_13](https://doi.org/10.1007/978-1-4020-6029-8_13)
- Al-Noor S. Population Status of Gold Fish *Carassius auratus* in Restored East Hammar Marsh, Southern Iraq. Journal of King Abdulaziz University-Marine Sciences. 2010;21(1):65–83.

- Available: <https://doi.org/10.4197/Mar.21-1.3>
8. Arindam M, Paramveer S, Manas M, Singh M, Girish T, Gaurav, Tripathi. Comparative study of gold fish (*Carassius auratus*) breeding via induced and natural breeding. Unpublished; 2018.  
Available:  
<https://doi.org/10.13140/RG.2.2.15956.04484>
  9. Chanda M, Paul M, Maity J, Dash G, Gupta S, Patra B. Ornamental fish goldfish, *Carassius auratus* and related parasites in three districts of West Bengal, India. *Chronicles of Young Scientists*. 2011;2(1):51.  
Available:<https://doi.org/10.4103/2229-5186.79351>
  10. Faizullah M. Impact of biofloc technology on the growth of goldfish young ones. *Indian Journal of Science and Technology*. 2015;8(1):1–8.  
Available:<https://doi.org/10.17485/ijst/2015/v8i13/54060>
  11. Patil PA, Dube K, Verma AK, Chadha NK, Sundaray JK, Jayasankar P. Growth performance of goldfish, *Carassius auratus* and basil, *Ocimum basilicum* in media bed aquaponics. *Indian Journal of Fisheries*. 2019;66(1).  
Available:<https://doi.org/10.21077/ijf.2019.66.1.78353-15>
  12. Jitendrasinh, Rana Rushirajsinh, Kotiya AS, Jungivala Mansi Dipakbhai. "Bioactive Feed Ingredients Used in Aquaculture: A Review". *Journal of Scientific Research and Reports*. 2024;30 (5):399-414.  
Available:<https://doi.org/10.9734/jsrr/2024/v30i51956>.
  13. Mangiamele LA, Thompson RR. Testosterone rapidly increases ejaculate volume and sperm density in competitively breeding goldfish through an estrogenic membrane receptor mechanism. *Hormones and Behavior*. 2012 Jul 1;62(2): 107-12.
  14. Tacon AG, De Silva SS. Feed preparation and feed management strategies within semi-intensive fish farming systems in the tropics. *Aquaculture*. 1997 May 15;151(1-4):379-404.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/119519>