



Tomato Performance in a Protected Structure: A Review

Aqleema Banoo ^{a*}, Shabber Hussain ^a, Nazir Hussain ^b,
Anwar Hussain ^c, F. A. Shah Khan ^a
and Shahnawaz Rasool Dar ^d

^a KVK Kargil-II Zanskar, SKUAST-K, J&K, India.

^b KVK Kargil-I Kargil, SKUAST-K, J&K, India.

^c High Mountain Arid Agriculture Research Institute Leh, UT Ladakh, India.

^d Research Center for Residue and Quality Analysis, SKUAST-K, J&K, India.

Authors' contributions

"This work was carried out in collaboration among all authors. Author AB and SH designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors FASK and SRD managed the analyses of the study. Author NH and AH managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/air/2024/v25i51134>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121037>

Review Article

Received: 14/06/2024

Accepted: 16/08/2024

Published: 24/08/2024

ABSTRACT

In the event of climate change, sheltered agriculture might offer plenty of opportunity for high-value vegetable crop production to occur during the "off" season. Tomato (*Solanum lycopersicum* L.) is one of the most significant high-value crops and a rich source of lycopene, β -carotene, phenols, flavonoids, and several vitamins. Growers can produce tomato fruit during the off-season and get higher prices in the market by using protected buildings like net houses, polyhouses, semi-high tunnels, and mulching. Through the modification of the natural environment, this technology has significantly improved productivity for areas with unfavourable weather conditions. In addition, it has

*Corresponding author: E-mail: aqleemabanoo786@gmail.com;

Cite as: Banoo, Aqleema, Shabber Hussain, Nazir Hussain, Anwar Hussain, F. A. Shah Khan, and Shahnawaz Rasool Dar. 2024. "Tomato Performance in a Protected Structure: A Review". *Advances in Research* 25 (5):29-37. <https://doi.org/10.9734/air/2024/v25i51134>.

extended the harvest period, increased yields, improved quality, increased production stability, and made fresh, high-quality produce available during the off-season, especially in Himalayan areas. From this study, it is pertinent to opine that tomato is high value vegetable crop in which protected cultivation is always advantage over open field cultivation. We highlight or briefly discussed the differences between growing tomatoes in an open field and under protection structure, as well as the growth characteristics, yield, and other quality parameters in this review.

Keywords: *Tomato; Solanum lycopersicum; protected cultivation; open field condition; yield.*

1. INTRODUCTION

Tomato (*Solanum lycopersicum* L.) is one of the most widely-grown greenhouse vegetables in the world [1], and is highly demanded by fresh market. It plays a vital role in human health also [2]. "Fresh tomato production was around 180 million tonnes worldwide in 2020. The top five tomato-producing nations are China, India, Pakistan, Turkey, and the United States, together generating more than 60% of the world's tomatoes. China is the country with the largest tomato production in the world, with an annual output of more than 65.15 million tonnes, accounting for one-third of the global tomato production" [3]. "It's output in India reached a peak in 2019–20 at 21.187 million tonnes (MT), and it has subsequently been in decline. It decreased to 20.69MT in 2021–2022 and 20.62MT in 2022–2023" [4]. "Tomatoes are high in vitamin C, potassium, folate, and vitamin K, all of which are responsible for a variety of health benefits, including cancer prevention, heart health, and constipation prevention" [5], [6], [7]. "Furthermore, tomatoes have significant nutritional value and positive health impacts due to their potassium, ascorbic acid (Vitamin C), and lycopene levels. Tomato continues to be the most important vegetable in the world due to increasing commercial and dietary value, widespread production as well as model plant for research purpose" [8]. Similarly, tomato is utilized as a fresh crop or processed into various forms such as paste, puree and juices.

"Tomatoes are planted both in the greenhouse and open fields. Therefore, developing a sustainable production system is essential for improved crop growth and a consistently high tomato yield" [3]. "In the plains of India, tomato is mainly grown as Rabi crop. However, in the Himalayan region, it is mostly grown as summer season crop. In in trans-Himalayan region of Ladakh its production is limited due to harsh climate, short growing season and extreme temperature fluctuations. Hence, protected cultivation is a feasible answer for successful

cultivation of tomato in this region" [9]. "With the innovative technique of protected cultivation, environmental conditions can be partially or completely controlled" [10], [11], [12]. Singh and Asrey [13] "recommended that cultivation of tomato in a greenhouse would help to obtain high productivity and better return. The greenhouse environment encompasses climatic and nutritional elements, along with structural and mechanical conditions". "If properly managed, it can significantly increase fruit quality and yield. Since the costs associated with greenhouse cultivation are generally higher, due to the initial investments in the structure" [14]. The protected cultivation of tomato is always aimed to obtain high yield, good quality crop, improved shelf life and year around availability of tomato fruit [15], [16], [17]. "The parameters of economic importance like fruit weight and yield are better under protected conditions but the nutritional parameters like vitamin C and lycopene content are higher in open conditions" [18]. "Cultivation of tomatoes under protected structure protects tomato fruit from biotic and abiotic stresses especially, during the off-season. Also increases yield by increasing fruits per plant, fruit weight, fruit length and fruit diameter" [19], [15]. "Under protected structures such as net houses and polyhouses, growers can produce tomato fruit during the off-season and receives higher prices in the market. If these structures are carefully maintained, it will increase productivity and net returns per unit area, and also keep pests away" [20]. Rana et al. [18] "also observed that tomato crop grown extensively in the plastic greenhouses increase productivity". "The environmental factors affecting greenhouse tomato include air and root-zone temperatures, relative humidity (RH), light conditions (L), disease and insect intervention, as well as carbon dioxide" [21]. "Microclimatic parameters are directly affected the fruit yield and quality of tomato" [22], [14]. "Different techniques and methods of determinant and indeterminate varieties of tomato planting under open-field and greenhouse conditions are available in the textbooks" of [23], [24], [25], [21].

The main obstacles to growing vegetables particularly tomato is climate change, shrinking agricultural land, water shortages, rising seed and input costs, a lack of qualified labour, market price volatility, the appearance of illnesses and insect resurgence, etc. In order to support the protected production of tomatoes and to recommend various scholars for upcoming research and development, this review paper has been painstakingly documented.

2. PERFORMANCE OF TOMATO UNDER PROTECTED CULTIVATION AS COMPARE TO OPEN FIELD CULTIVATION

Protected cultivation of vegetables endorses effective use of the soil nutrients, soil moisture, humidity, temperature, CO₂, light intensity etc., which results in increase the production and quality vegetables by delimiting biotic and abiotic stresses whereas in open cultivation plants suffer a great damage due to weeds, insects and diseases with limited use efficiency of natural resources and environment factors. Greenhouse is the present trends in Indian agriculture where the crop is raised under fully automated or partly artificially manipulated environment endorse higher production and productivity as compared to open filed cultivation [11]. Takteet et al. [26] revealed that polythene sheet and shading nets avoids extreme sunshine, chilling, freezing, high wind and biotic stresses. Environmental factors like day and night temperature, air relative humidity, light intensity, light period, CO₂ concentration, water availability, nutrient availability, plant population, plant media etc. are significantly maintained in greenhouse for crop cultivation [10], [18]. There is drastic increase in the resource-use efficiency through protected crop cultivation in comparison to the conventional open field cultivation [27]. Tomato production has undergone many changes in the way as it is being grown in different regions, since different limiting factors including climatic conditions, availability of water and nutrients, and quality and quantity of light must be taken into account to provide appropriate conditions for the optimal growth and development of the crop. Because of their sophisticated light-perception system, plants are able to modify and optimize their metabolism and performance in response to the specific environmental factors that are most common. As observed by Arsovski et al. [28], that light affects many developmental and physiological processes which include germination [29], flowering [30] and direction of growth [31]. The

growing conditions especially light and temperature are also known to influence both composition and quality of tomato fruits. [32] “reported that plant height is better in polyhouses as compared to open field conditions”. “The environment in the polyhouse favour the growth and development of tomato plant through increased plant height, number of branches, stem grith and yield” [33], [34], [18]. “However, fruits harvested from open field had higher TSS, acidity, total sugar, ascorbic acid and the lycopene content than the fruits grown under protected conditions” [35], [18]. “Therefore, both types of environmental conditions are favourable in some aspects. Protected cultivation had a significant impact in improving the crop productivity, employment and income generation. Furthermore, protected cultivation attains more stable income than open field cultivation” [36], [37], [38], [39]. Therefore, it is necessary to grow tomatoes in protected environments like net houses, poly houses, and green houses, among others, in order to acquire high-quality produce and production throughout the off-season. Green houses have a great deal of potential to increase tomato yield [40], [41].

3. PERFORMANCE OF CAPSICUM FOR VEGETATIVE GROWTH PARAMETERS

“An appropriate environmental condition inside polyhouse promotes robust cell division and cell enlargement in the terminal portion of the plant, which supports the maximum plant height, the greater number of leaves per plant, a greater number of branches per plant, and overall plant growth. In Tomato, a greater number of branches and better plant height reported in polyhouses as compared to open field conditions” [32], [9]. “Total fruit bearing period was also prolonged under polyhouse. For that reason, total number of fruit harvests was more in polyhouse than open field” [42], [43]. Plant spacing and stem pruning are the two most important factors. Optimum plant spacing may help in proper utilization of land and high-quality fruits. However, stem pruning may influence the production of yield [44]. Suitable day and night temperature, relative humidity and light intensity are available in polyhouse resulting in increasing yield by improving the vegetative characters in tomato [45]. Foolad and Lin [46] attempted to connect tomato vegetative development to cold tolerance during seed germination. They discovered that there was no connection between the two and suggested that distinct

genes are engaged at these developmental stages. Therefore, it's possible that characteristics that affect fruit output at lower temperatures have little to do with growth during the vegetative stage. Ulukapi et al. [47] reported that the average fruit weight and quality increased with fruit reduction on the stem, and accordingly the plant yield increased. Leaf and side branch pruning has been shown to improve final output in a number of ways. According to Khoshkam et al. [48], "training increases a plant's capacity to absorb sunlight, which is essential for its growth and development. It is also important to maintain adequate air movement around the plant to reduce risk of fungus and insect problems. A dense canopy of leaves shades the fruits, causing them to be pale". Rana *et al.* [18] "reported that tomato plants grown in polyhouse climate produced about 50% higher fruit yield than the tomato plants grown in open field conditions". "The significantly higher yield in the plants grown under polyhouse condition over the plants grown in the open field was associated with the production of higher number of fruits with greater length and diameter and fruit weight than those in open field" [34], [15].

Dixit et al. [41] "reported that different combinations of organic manures i.e., FYM, Vermicompost and Poultry manure are the best treatment for better plant growth and flowering of tomato under protected structure". Reddy et al. [49], Patil et al. [50], Poul et al. [51] "also reported that organic manure increase plant growth and yield". "The use of FYM was found to be the most important variable and showed a significantly positive relationship with the yield of tomato under-protected and open-field conditions as reported" by [39].

4. PERFORMANCE OF TOMATO FOR YIELD PARAMETERS

Tomato yield is a function of the entire plant's growth rather than a single feature. A tomato plant will never produce a large amount of yield if it does not grow well. Thus, the relationship between plant form, physiology, and growth circumstances determines yield [23]. "Tomato yield in the tropics is much lower than that in the temperate zone due to several factors like high humidity, high temperature, excessive rainfall, disease and pest" [52]. Temperature has a major impact on crop phenological growth and output [53]. Plants grown in greenhouse have favourable environmental conditions that enable more effective use of water and nutrients,

ensuring that the crop's genetic potential is fully expressed for growth and production. However, decrease in temperature will also affect the growth and development of plant including biomass production, fruit growth and yield [54], [23]. According to Adams et al. [45], "trusses form more quickly at higher temperatures. Because of the shorter fruit growth period and faster initiation of new trusses at higher temperatures, early yields are higher" [55], [45]. "Tomato plant grown under polyhouse was observed to be earlier in flowering and fruit setting by about 3 and 8 days, respectively when compared to the crop raised under open condition" [32]. "Microclimate inside protected structures influences the flowering and fruiting during winter months which is helpful for higher yield in tomato" [56], [34], [15]. Kumar and Arumugam [57] "observed that the growth and yield parameters of different vegetables were significantly enhanced under protected over open field". "Similar results were also observed" by [58]. Parvej et al. [15] "reported that under polyhouse plants had higher number of flower clusters/plant, flowers/cluster, flowers/plant, fruit clusters/plant, fruits/cluster and fruits/plant, and fruit length, fruit diameter, individual fruit weight, fruit weight/plant and fruit yield than open field condition. Kanwar [9] "reported that variety 'Shivalik' performed best with respect to yield followed by 'Pusa Rohini' under polyhouse conditions. Moreover, cultivation of tomato under the polyhouse produced 136.12% more yield per ha and 188.93% more fruits per plant compared to open field cultivation". Khadka and Adhikari [59] "reported that not only yield but also gross return, gross margin and net profit from polyhouse farming of tomato was 60.14%, 68.25%, 74.32% and 69.97% respectively higher than that of open field farming". Kanthaswamy et al. [60], Nagalakshmi et al. [61], Gaikwad et al. [62], Srivastava et al. [63], Mishra et al. [33], Kang and Sidhu [34] "reported that higher values of all the yield components and yield of tomato crop grown under polyhouse was due to the taller plants and much number of branches with greater Leaf area index attributed by warmer temperature". "Although lower amount of available PAR (Photosynthetically active radiation) under polyhouse could not affect the growth and yield of tomato" [19], [64]. "Overall, to obtain higher yields under this production system, it is important to reduce the elevated temperature and increase the relative humidity inside the protective structures to be adapted for local growing conditions. The improved microclimate within the polyhouse, which

included winter temperatures 4-9°C higher than in the adjacent open field, was primarily responsible for the early and higher output of several vegetable crops” [65]. Thus, the polyhouse environment may provide a new scope for commercial production of high value vegetable crops like tomato.

5. PERFORMANCE OF TOMATO FOR QUALITY PARAMETERS

Plants grown in open conditions face unfavourable weather and biotic stresses, which leads to lower-quality yield production. In contrast, plants grown in greenhouses yield higher-quality produce because of favourable environmental conditions and lower rates of weeds, diseases, and pests. “Morphological development like plant height, number of branches per tomato plant, leaf area expansion rate and leaf area index were positively favoured due to the warmer environment inside the polyhouse” [66], [67], [42]. “Environmental factors, including light and temperature, may influence the chemical and nutritional composition of tomatoes [68-77], including lycopene, accounting for 80-90% of the total pigments presented in ripened tomato” [78]. Hormonal action is influenced by environmental conditions, which can lead to early blooming, fruit setting, increased fruit production, and increased fruit count. Occurrence of various limiting factors like biotic and abiotic stresses can be minimized under protected structures [17]. Moreover, fruits produced in open field were more acidic (0.49%) than fruits produced in a protected environment (0.40%) [18]. Frusciante et al. [79] “reported that fruit grown open filed condition have higher ascorbic acid rather than fruit grown in protected environment condition. It may be a result of the lower photosynthetic activity of the plant (shading in protected environment) in this environment and lower carbohydrate accumulation in the fruits”. Rana et al. [18] “also observed that Genotype influenced the lycopene content of fruit, while the environment did not influence this characteristic”. “Cherry tomatoes have also been earlier reported to contain higher content of lycopene than local ones under open conditions” [70].

“Moreover, studies had shown that color shading structures improved tomato fruit quality and carotenoid content” [17]. “Because coloured shade nets enable a better use of the sun’s energy due to the modulation of the radiation spectra that reach the crops, thus promoting

physiological responses in plant and fruit development, including leaf area index, chlorophyll and carotenoid contents, tissue structure, fruit ripening, physiological disorders, nutritional quality, etc” [71]. “In contrast, during tomato production, high temperatures and high solar radiation may reduce lycopene and β -carotene levels resulting in fruit damage, sunscald and increasing unmarketable fruit yield” [72]. Studies also reported that colour net houses provide a positive effect on tomato production in terms of improvement in morphometric parameters like taller plants, larger terminal leaflets compared to those plants grown under open field conditions. According to [16] “tomato varieties grown in open fields and poly houses have varying in flowering time. Phenological development governs the plant growth and productivity”. “Days to flowering, fruiting and maturity of crop are the important phenological events which determine the productivity of a crop” [15].

6. CONCLUSION

From this study, it is pertinent to opine that tomato is high value vegetable crop in which protected cultivation is always advantage over open field cultivation. Naturally ventilated greenhouse cultivation endorses maximum plant height, number of leaves, number of branches per plant and overall vegetative growth in tomato. Moreover, Improved yield parameters such as early flowering and fruiting, a greater number of flower and fruits per plant, better fruit set, early fruit ripening, maximum fruit size and weight, early first harvesting, total yield are increased in tomato with superior quality fruit are obtained under polyhouse growing.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENT

Authors take this opportunity to express our deep sense of gratitude and sincere thanks to both the organization (Funding agency DST SEED and administrative body KVK).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Viuda-Martos M, Sanchez-Zapata E, Sayas-Barbera E, Sendra E, Perez-Alvarez JA, Fernandez-Lopez J. Tomato and tomato by products. Human health benefits of lycopene and its application to meat products: A review. *Critical Reviews in Food Science and Nutrition*. 2014; 54(8):1032-1049.
2. Li N, Wu X, Zhuang W, Xia L, Chen Y, Wu C, Rao Z, Du L, Zhao R, Yi M, Wan Q. Tomato and lycopene and multiple health outcomes: Umbrella review. *Food Chemistry*. 2021;343:128396.
3. Zhang F, Lv Z, Zhang H, Guo J, Wang J, Lu T, Zhangzhong L. Verification of improved yolox model in detection of greenhouse crop organs: Considering tomato as example. *Computers and Electronics in Agriculture*. 2023; 205:107582.
4. FAO. FAOSTAT. Available:<http://www.fao.org/faostat/en/home>. Accessed 18 Jan 2021.
5. Megan LD. Everything you need to know about tomatoes. 2017. Available:<https://www.medicalnewstoday.com/articles/273031>. Accessed 27 Dec 2020
6. Ramasamy S, Ravishankar M. Integrated pest management strategies for tomato under protected structures. In: *Sustainable Management of Arthropod Pests of Tomato*. 2008;2:313-22.
7. Islam J, Kabir Y. Effects and mechanisms of antioxidant rich functional beverages on disease prevention. In: *Functional and Medicinal Beverages*. 2019;11:157-198.
8. Kimura S, Sinha N. *Tomato (Solanum lycopersicum): A model fruit-bearing crop*. Cold Spring Harbor Protocols. 2008.
9. Kanwar MS. Performance of tomato under greenhouse and open field condition in the trans- Himalayan region of India. *Advance in Horticulture Science*. 2011;25(1):65-68.
10. Baghel BS, Gupta M, Tiwari R. Potential and prospects of protective cultivation of horticultural crops. *Proc. All India Sem. on Potential and Prospects for Protective Cultivation organized by the Institute of Engineers*. 12 December. Ahmednagar. 2003;12-13.
11. Navale AV, Nandagude SB, Pawar AG, Ghodke HM, Bhosale AD. Comparative study of skirting and top covering effect in low-cost greenhouse. *Proceeding of All India on Seminar Potential and Prospects for Protective Cultivation organized by the Institute of Engineers*. December. Ahmednagar. 2003;12-13.
12. Wani KP, Singh PK, Narayan N, Khan SH, Amin A. Prospects of vegetable production in cold arid region of Ladakh, Achievement and future strategies. *International Journal of Current Research*. 2011;33(6):10-7.
13. Singh R, Asrey R. performance of tomato and sweet pepper under unheated greenhouse. *Haryana Journal of Horticulture Science*. 2005;34(1&2):174-175.
14. Shamshiri RR, Jones JW, Thorp KR, Ahmad D, Man HC, Taheri S. Review of optimum temperature humidity and vapour pressure deficit for microclimate evaluation and control in greenhouse cultivation of tomato: A review. *International Agrophysics*. 2018;32(2):287-302.
15. Parvej MR, Khan MAH, Awal MA. Phenological development and production potentials of tomato under polyhouse climate. *Journal of Agricultural Sciences–Sri Lanka*. 2010;5(1).
16. Gupta T, Thapa U, Kumar P. Assesment of Variety of Tomato for Growth Parameters under Cover and Open Field Conditon. *International Journal of Pure & Applied Bioscience*. 2017;5(6):1238-1245.
17. Sotelo-Cardona P, Lin MY, Srinivasan R. Growing tomato under protected cultivation conditions: Overall effects on productivity nutritional yield and pest incidences. *Crops*. 2021;1:97–110.
18. Rana N, Kumar M, Walia A, Sharma S. Tomato fruit quality under protected environment and open field conditions. *International Journal of Bio-resource and Stress Management* 2014;5(3): 22-426.
19. Aberkain K, Gosselin A, Vineberg S, Dorais M. Effects of insulating foams between double polyethylene films on light transmission growth and productivity of greenhouse tomato plants grown under supplemental lighting. *Acta Horticulturæ (ISHS)*. 2006;711:449-454.
20. Ramasamy S, Ravishankar M. Integrated Pest Management Strategies for Tomato Under Protected Structures. In: *Sustainable Management of Arthropod Pests of Tomato*. Academic Press. 2018; 313-322.
21. Jones JB. *Instructions for growing tomatoes in the garden and green-house*. Gro Systems Anderson SC USA; 2013.

22. Kittas C, Katsoulas N, Rigakis V, Bartzanas T, Kitta E. Effects on microclimate crop production and quality of a tomato crop grown under shade nets. *The Journal of Horticultural Science and Biotechnology*. 2012;87(1):7-12.
23. Van der Ploeg A, Heuvelink E. Influence of sub-optimal temperature on tomato growth and yield: a review. *Journal of Horticultural Science & Biotechnology*. 2005;80(6):652-659.
24. Cherie E. *The complete guide to growing tomatoes: A complete step-by-step guide including heirloom tomatoes (back-to-basics gardening)*. Atlantic Publishing Group Inc., Ocala, Florida. 2010.
25. Hochmuth GJ, Hochmuth RC. *Production of greenhouse tomatoes - Florida greenhouse vegetable production handbook Selection of Cultivars. Production*. 2012;3:1-18.
26. Takte RL, Ambad SN, Kadam US, Dhawale BC. Greenhouse cladding material shade nets ventilation etc. *Proceeding of All India on Sem Potential and Prospects for Protective Cultivation Organized by the Institute of Engineers. Ahmednagar*. 2003;117-119.
27. Choudhary AK. Scaling-up of protected cultivation in Himachal Pradesh India. *Current Science*. 2016;111(2):272-277.
28. Arsovski AA, Galstyan A, Guseman JM, Nemhauser JL. *Photomorphogenesis. The Arabidopsis Book. American Society of Plant Biologists*. 2012;10.
29. Bentsink L, Koornneef M. *Seed Dormancy and Germination. The Arabidopsis Book. American Society of Plant Biologists*. 2008;6.
30. Alvarez-Buylla ER, Benitez M, Corvera-Poire A, Chaos-Cador A, de Folter S, Gamboa de Buen A, Garay-Arroyo A, Garcia-Ponce B, Jaimes-Miranda F, Perez-Ruiz RV. *Flower Development. The Arabidopsis Book. American Society of Plant Biologists*. 2010;8.
31. Pedmale UV, Celaya RB, Liscum E. *Phototropism: Mechanism and Outcomes. The Arabidopsis Book. American Society of Plant Biologists*. 2010;8.
32. Ganesan M. Effect of poly-greenhouse models on plant growth and yield of tomato (*Lycopersicon esculentum*). *Indian Journal of Agricultural Sciences*. 2002;72(10):586-588.
33. Mishra JN, Molianty BK, Pradhan PC, Naik P. Study on biometric characteristics of okra in greenhouse. *Orissa Journal of Horticulture*. 2003;31(1):112-113.
34. Kang BS, Sidhu BS. Studies on growing off-season tomato nursery under polyhouse. *Annals Agricultural Biology Research*. 2005;10(1):53-56.
35. Loures JL. *Estabelecimento e avaliacao do sistema de producao denominado Fito em estufa e campo. Ph.D. Thesis. Universidade Federal de Vicosa Vicosa Brazil*. 2001;109.
36. Singh B, Sirohi NPS. Protected cultivation of vegetables in India: Problems and future prospects. *Acta Horticulture*. 2006; 710: 339-42.
37. Singh AK, Singh B, Gupta R. Performance of sweet pepper (*Capsicum annum*) varieties and economics under protected and open field conditions in Uttarakhand. *Indian Journal of Agricultural Sciences*. 2011;81:973-975.
38. Mwangi WJ. *Comparative analysis of greenhouse versus open-field small scale tomato production in Nakuru-North district Kenya. Thesis M.Sc. Department of Agricultural Economics and Agribusiness Management Egerton University Kenya*; 2012.
39. Mehta K, Guleria JS, Thakur RK. Comparative Economics of Tomato Cultivation under Protected versus Open-field Conditions in Mid-hills of Himachal Pradesh. *Indian Journal of Economics and Development*. 2020;16(1):104-110.
40. Chandra P, Sirohi PS, Behera TK, Singh A K. *Cultivating vegetable in polyhouse. Indian Horticulture*. 2000;45:17-32.
41. Dixit PS, Prasad VM, Kasera S, Singh S. Effect of various organic inputs on growth and its attributes of tomato (*Solanum lycopersicum* L.) cv. GS-600 under polyhouse condition. *International Journal of Conservation Science*. 2018;6(6):2537-2539.
42. Pandey VK, Dwivedi SK, Pandey A, Sharma HG. Low cost polyhouse technology for vegetable cultivation in Chhattisgarh Region. *Plant Archives*. 2004;4(2):295-301.
43. Cargnelutti Filho A, Radin B, Matzenauer R, Storck L. Number of harvest and comparison of tomato genotypes cultivated under plastic greenhouse. *Pesquisa Agropecuaria Brasileira* 2004;39(10):953-959.
44. Ara N, Bashar MK, Begum S, Kakon SS. Effect of spacing and stem pruning on the

- growth and yield of tomato. International Journal of Sustainable Crop Production. 2007;2(3):35-39.
45. Adams SR, Cockshull KE, Cave CRJ. Effect of temperature on the growth and development of tomato fruits. Annals of Botany. 2001;88:869-877.
 46. Foolad MR, Lin GY. Relationship between cold tolerance during seed germination and vegetative growth in tomato: Germplasm evaluation. Journal of the American Society for Horticultural Science. 2000;125:679–683.
 47. Ulukapi K, Ercan N, Onus AN. Effects of Different Training Systems and Planting Density Combinations on Yield and Quality Parameters of Cherry Tomato Cultivar F1 M25. In IV Balkan Symposium on Vegetables and Potatoes. Acta Horticulturae. 2009;830:441-446.
 48. Khoshkam S, Seyedi Z, Aeen A. The impact of different plant training systems on quantitative and qualitative parameters of greenhouse tomato cultivars. International Journal of Farming and Allied Sciences. 2014;3(6):659-663.
 49. Reddy CS, Narayanamma M, Chiranjeevi C and Reddy IP. Effect of nutrient sources on the fruit yield of tomato (*Lycopersicon esculentum* Mill.). Vegetable Science. 2002;29(2):193-194.
 50. Patil MB, Mohammed RG, Ghadge PM. Effect of organic and inorganic fertilizers on growth yield and quality of Tomato. Journal of Maharashtra Agriculture University 2004;29(2):124-127.
 51. Poul AS, More SD, Lohit VD, Bodke RG. Effect of organic and inorganic nutrient sources on growth yield and nutrient uptake in Tomato. Journal of Soils and Crops. 2004;14(1):40-45.
 52. Opena RT. Development of Tomato and Chinese Cabbage cultivars adapted to the Hot Hamid tropics Acta Horticulture. 1985;153:421-436.
 53. Awal MA, Ikeda T, Itoh R. The effect of soil temperature on source–sink economy in peanut (*Arachis hypogaea*). Environmental and Experimental Botany. 2003;50(1):41-50.
 54. Heuvelink E. Effect of temperature on biomass allocation in tomato (*Lycopersicon esculentum*). Physiologia Plantarum. 1995;94:447–452.
 55. De Koning ANM. The effect of temperature on fruit growth and fruit load of tomato. Acta Horticulturae. 1989;248:329–337.
 56. Rylski I, Aloni B, Karni L and Zaidman Z. Flowering fruit set fruit development and fruit quality under different environmental conditions in tomato and pepper crops. Acta Horticulture. 1994;366:45–56.
 57. Kumar RS, Arumugam T. Performance of vegetables under naturally ventilated polyhouse conditions. Mysore Journal of Agricultural Sciences. 2010;44:770-776.
 58. Kumar P, Chauhan RS, Grover RK. Economics analysis of tomato cultivation under poly house and open field conditions in Haryana India. Journal of Applied and Natural Science. 2016;8(2):846-848.
 59. Khadka S, Adhikari RK. Comparative economics of tomato production under polyhouse and open field condition in Dhading district of Nepal. 2012; 100-107.
 60. Kanthaswamy V, Narendra S, Veeraragavathatham D, Srinivasan K, Thiruvudainambi S. Studies on growth and yield of cucumber and sprouting broccoli under polyhouse condition. South Indian Horticulture. 2000;48(1/6):47-52.
 61. Nagalakshmi S, Nandakumar N, Palanisamy D, Sreenarayanan VV. Naturally ventilated polyhouse for vegetable cultivation. South Indian Horticulture. 2001;49:345-346.
 62. Gaikwad AM, Katwate SM, Nimbalkar CA. Evaluation of chrysanthemum varieties under polyhouse conditions. South Indian Horticulture. 2000;50(4/6): 624 628.
 63. Srivastava P, Srivastava BK, Singh MP. Effect of date of planting and growing environment on the plant survival, growth and yield of early cauliflower in rainy season. Vegetable Science. 2000;29(2): 157-160.
 64. Marcelis LFM, Broekhuijsen AGM, Meinen F, Nijs FMFM, Raaphorst MGM. Quantification of the growth response to light quantity of greenhouse-grown crops. Acta Horticulturae (ISHS). 2006;711:97-103.
 65. Cheema DS, Kaur P, Kaur S. Off-season cultivation of tomato under net house conditions. Acta Horticulturae. 2004;659:177-181.
 66. Duhr E, Dubas A. Effect shading the soil with plastic film on the dynamics of plant development and yield of maize sown on different dates. Prce Kumisji Nouk Rolnicsy Ch-I-Kamiji-lesnych. 1990;69:9-18.
 67. Miah MM. Performance of five winter vegetables under different light conditions

- for Agroforestry systems. M. S. Thesis, BSMRAU, Gazipur, Bangladesh; 2001.
68. Garcia-Valverde V, Navarro-Gonzales I, Garcia-Alonso J, Periago JM. Antioxidant bioactive compounds in selected industrial processing and fresh consumption tomato cultivars. *Food Bioproduction Technology* 2013;6:391–402.
69. Frusciante L, Barone B, Carputo D, Ercolano MR, Della Roca F, Esposito S. Evaluation and use of plant biodiversity for food and pharmaceuticals. *Fitoterapia* 2000;71:66-72.
70. Raffo A, La Malfa G, Fogliano V, Maiani G and Quaglia G. Seasonal variations in antioxidant components of cherry tomatoes (*Lycopersicon esculentum* cv. Naomi F1). *Journal of Food Composition and Analysis* 2006;19:11-19.
71. Ilić ZS, Fallik E. Light quality manipulation improves vegetable quality at harvest and postharvest: A review. *Environmental and Experimental Botany*. 2017;139:79-90.
72. Diaz-Perez JC. Bell pepper (*Capsicum annuum* L.) crop as affected by shade level: Fruit yield quality and postharvest attributes and incidence of phytophthora blight (caused by *Phytophthora capsici* Leon.). *HortScience*. 2014;49:891-900.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/121037>