



Assessment of Nutritional Status among Male and Female Undergraduate Students of Saveetha Dental College

M. Preethi Raj^a, A. Jothi Priya^{b*†} and R. Gayatri Devi^{b†}

^a *Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai - 600077 India.*

^b *Department of Physiology, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai - 600077 India.*

Authors' contributions

This work was carried out in collaboration among all authors. Author AJP performed the statistics approval, approval of the manuscript. Author MPR managed the Literature search, data collection, analysis and wrote first draft of the manuscript. Author RGD approved the final manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i60B34963

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/78822>

Original Research Article

Received 20 October 2021
Accepted 25 December 2021
Published 27 December 2021

ABSTRACT

A rational way of nourishment, combined with adequate physical activity, are the basic components of maintaining proper body condition. The aim of the study was to evaluate nutritional status among male and female undergraduate students of Saveetha dental college. A cross sectional survey of different questions, dietary patterns was conducted with a sample of 184 students (I, II, III, IV, and V students). Survey was conducted via google form through random sampling to identify undernutrition in the sample. Questionnaires were based on dietary health patterns and nutritional status in day to day life. The sample size for this study is about 184 participants and at the end of the survey, all the data were collected and analysed by using IBM SPSS software. The results obtained was the majority of the students are at normal weight and the scores met optimal levels. This suggests good quality and quantity of diet.

[†]Assistant Professor;

*Corresponding author: E-mail: jothipriya.sdc@saveetha.com, a.jothipriya88@gmail.com;

Keywords: Nutritional status; health; diet; malnutrition; questionnaires; IBM SPSS software.

1. INTRODUCTION

The major determinants of health and non communicable diseases are nutritional status and diet [1]. One of the importance of public health is nutritional status. This attracted the attention of economists and other social activists [2,3,4]. Students entering university environments have new social relations and behaviours - eating habits, physical activity-smoking, alcohol consumption etc [5]. This interferes in social, physical and biological problems leading to health problems. The transition to adolescence is an important period for change in behavioural patterns which has a risk factor to chronic disease [6].

It is very important to recognise and treat malnutrition because it's consequences include improper functioning of respiratory muscles and renal functions, the contractility of cardiac muscles gets reduced [7]. Nowadays, in developing countries one of the serious health problems and highly prevalent is undernutrition [8]. This occurs due to the intake of net nutrients is less than requirements [9]. It is easy to reach children through institutions such as schools and colleges. We have the responsibility for society to do our best to protect young people [10]. College is a critical period where lifelong lifestyle habits are developed, which have a lasting impact on development of chronic diseases [11,12,13].

Nutrition education has a profound impact on dietary habits and food choices in many college students [14,3]. Thus, schools and colleges represent an opportunity for students to learn new skills and lead healthy lifestyle practices. Students will have necessary knowledge and skills which are essential to make healthy lifestyle choices [15,12]. During college years, women face a great pressure to reduce their weight to achieve the 'thin ideal' body. Being thin makes women more attractive to their peers and more likely to gain peers' attention [16,17]. Students need to be healthy, physically active, and well nourished in order to succeed in their academic studies [16,18].

Different factors are associated with malnutrition among undergraduates of Saveetha Dental College. One set of factors refers to demographic details such as age, gender and other questionnaires [19]. Another set of factors consists of college, year of study etc., [15]. This

research conducted on the topic of assessment of nutritional status among male and female of Saveetha dental college has focused on undergraduate students to determine the underweight or malnutrition groups who are at risk of nutrient deficiencies [20]. These kinds of surveys and studies will be useful for guiding, planning and also implementing health interventions [21].

2. MATERIALS AND METHODS

A cross-sectional survey was conducted among the dental students with a sample size of 184. A self administered structured questionnaire was prepared based on Nutritional Status among male and female undergraduate students of Saveetha Dental Students and consisted of 11 questions. It was circulated to the students through an online platform (google form). The statistics were done using SPSS software, and the chi-square test was done. The pros of the survey is that the students of various Institutions were surveyed. Simple random sampling method was the sampling method used to minimise the sampling bias.

3. RESULTS AND DISCUSSION

In our study we found that; 74.46% of the population are interested in healthy dietary patterns (Fig. 1). 48.37% of the population prefer Non vegetarian (Fig. 2). 67.93% of the population consume milk and milk products (Fig. 3). 71.74% of the population take snacks while watching television (Fig. 4). 44.02% of the population consume bakery products (Fig. 5). 25% of females (green colour) don't skip their meals which are considered to be statistically not significant (Fig. 6). 24.46% of females (blue colour) have a duration of 1 hour between their dinner and sleep. This shows statistically not significant (Fig. 7).

4. DISCUSSION

This study examined the knowledge of assessing the nutritional status among male and female undergraduate students of Saveetha dental college. According to the survey, it is observed that students responded are of normal weight and health dietary patterns but there is a slight difference in gender. That is, females are slightly weak compared to males. Compared to other research, the number of males participated was

264 and females was 300 but in this research, females were 100 whereas males were 84. In similar findings by 66% consume milk and milk products whereas 34% don't consume. In this study, 67.93% responded yes and 32.07% responded no. In the research by [22,23] 8% of males and 9% of females met the recommended

dietary intake of at least five servings of fruits and vegetables per day. In this study, 45% consume fruits everyday whereas 8.15% consume fruits in rare cases. The research done by [24] reported that females were more interested in changing their dietary habits and levels of physical activity than male students. In




Fig. 1. Pie chart showing percentage distribution of interest in health dietary patterns. Wherein the green colour represents yes (74.46%) and blue colour represents no (25.54%)




Fig. 2. Pie charts showing percentage distribution of food preference. Wherein blue represents both veg and non veg (25%), green colour represents non veg (48.37%) and beige colour represents vegetarian (26.63%)




Fig. 3. Pie chart showing percentage distribution of consumption of milk and milk products. Wherein blue represents no (32.07%) and green represents yes (67.93%)




Fig. 4. Pie chart showing percentage distribution of snacking while watching television. Wherein blue represents no (28.26%) and green represents yes (71.74%)

this study, 74.46% are interested in health dietary patterns whereas 25.54% are not interested. From this research, the limitations faced were time lagging and limited information. This establishes to promote nutritional status and dietary patterns among undergraduate students and also provides baseline data for further studies. Dietary health patterns, nutritional

status are the factors which play an important role in developing countries like India [25-29]. Very few of the undergraduate students who participated are overweight which leads them at risk. Most of the students are normal with proper health and BMI. They are well nourished [30-35].




Fig. 5. Pie chart showing percentage distribution of consumption of bakery products. Where, blue represents monthly once (23.91%), green colour represents once a week (32.07%) and beige colour represents twice a week (44.02%)




Fig. 6. Bar graph depicts the association between gender and skipping meals. X-axis represents the gender and the Y-axis represents the number of participants. Blue colour represents maybe, Green colour represents no and beige colour represents yes. 25% of females don't skip their meals, 22.83% of males don't skip their meals. This difference was statistically not significant (Pearson's chi square value: 2.281, p-value: 0.320, (p>0.05) - not significant)




Fig. 7. Bar graph depicts the association between gender and sleep duration. X-axis represents the gender and the Y-axis represents the number of participants. Blue colour denotes duration of 1 hour, Green colour represents duration of 2 hours and beige colour represents duration of 3 - 4 hours. 24.46% of females have 1 hour duration between their dinner and sleep, 20.11% of males have 1 hour duration between their dinner and sleep. This difference was statistically not significant (Pearson’s chi square value: 0.377, p-value: 0.828, ($p > 0.05$) - not significant)

5. CONCLUSION

Developing gender-specific programmes for promoting healthy lifestyle behaviours among students is recommended. In brief, this study provided preliminary data on various nutrition-related factors that can influence students’ health. Results can be useful for health educators for designing health promotion and nutrition education programmes aimed at improving students’ dietary and lifestyle behaviours. The data shown here is concluded by saying that both male and female students are of normal health. But a slight difference takes place in between them.

CONSENT AND ETHICAL APPROVAL

As per international standard or university standard guideline participant consent and

ethical approval has been collected and preserved by the authors.

ACKNOWLEDGEMENT

We thank all the participants and Saveetha Dental College for their support to conduct this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Abeywickrama HM, Swarna Wimalasiri KM, Koyama Y, Uchiyama M, Shimizu U, Chandrajith R, et al. Assessment of Nutritional Status and Dietary Pattern of a

- Rural Adult Population in Dry Zone, Sri Lanka [Internet]. International Journal of Environmental Research and Public Health. 2019;17:150.
Available:<http://dx.doi.org/10.3390/ijerph17010150>
2. Huda N, Ahmad R. Preliminary Survey on Nutritional Status among University Students at Malaysia [Internet]. Pakistan Journal of Nutrition. 2010;9:125–7.
Available:<http://dx.doi.org/10.3923/pjn.2010.125.127>
 3. Wadhwa R, Paudel KR, Chin LH, Hon CM, Madheswaran T, Gupta G, et al. Anti-inflammatory and anticancer activities of Naringenin-loaded liquid crystalline nanoparticles *in vitro*. J Food Biochem. 2021 Jan;45(1):e13572.
 4. Wahab PUA, Madhulaxmi M, Senthilnathan P, Muthusekhar MR, Vohra Y, Abhinav RP. Scalpel Versus Diathermy in Wound Healing After Mucosal Incisions: A Split-Mouth Study. J Oral Maxillofac Surg. 2018 Jun;76(6):1160–4.
 5. Barabadi H, Mojab F, Vahidi H, Marashi B, Talank N, Hosseini O, et al. Green synthesis, characterization, antibacterial and biofilm inhibitory activity of silver nanoparticles compared to commercial silver nanoparticles [Internet]. Inorganic Chemistry Communications. 2021;129:108647.
Available:<http://dx.doi.org/10.1016/j.inoche.2021.108647>
 6. Omege K, Omuemu VO. Assessment of dietary pattern and nutritional status of undergraduate students in a private university in southern Nigeria [Internet]. Food Science & Nutrition. 2018;6:1890–7.
Available:<http://dx.doi.org/10.1002/fsn3.759>
 7. Bharath B, Perinbam K, Devanesan S, AlSalhi MS, Saravanan M. Evaluation of the anticancer potential of Hexadecanoic acid from brown algae *Turbinaria ornata* on HT-29 colon cancer cells [Internet]. Journal of Molecular Structure. 2021;1235:130229.
Available:<http://dx.doi.org/10.1016/j.molstruc.2021.130229>
 8. Clarizia G, Bernardo P. Diverse Applications of Organic-Inorganic Nanocomposites: Emerging Research and Opportunities: Emerging Research and Opportunities. IGI Global. 2019;237.
 9. Puntis J. Nutritional assessment [Internet]. Oxford Medicine Online; 2018.
Available:<http://dx.doi.org/10.1093/med/9780198759928.003.0003>
 10. Fernández San Juan PM. Dietary habits and nutritional status of school aged children in Spain. Nutr Hosp. 2006 May;21(3):374–8.
 11. Contento IR, Koch PA, Lee H, Calabrese-Barton A. Adolescents demonstrate improvement in obesity risk behaviors after completion of choice, control & change, a curriculum addressing personal agency and autonomous motivation. J Am Diet Assoc. 2010 Dec;110(12):1830–9.
 12. Tahmasebi S, Qasim MT, Krivenkova MV, Zekiy AO, Thangavelu L, Aravindhan S, et al. The effects of oxygen-ozone therapy on regulatory T-cell responses in multiple sclerosis patients. Cell Biol Int. 2021 Jul;45(7):1498–509.
 13. Vivekanandhan K, Shanmugam P, Barabadi H, Arumugam V, Daniel Raj Daniel Paul Raj D, Sivasubramanian M, et al. Emerging Therapeutic Approaches to Combat COVID-19: Present Status and Future Perspectives. Front Mol Biosci. 2021 Mar 8;8:604447.
 14. Ha E-J, Caine-Bish N. Effect of nutrition intervention using a general nutrition course for promoting fruit and vegetable consumption among college students. J Nutr Educ Behav. 2009 Mar;41(2):103–9.
 15. Brown LB, Dresen RK, Eggett DL. College students can benefit by participating in a prepaid meal plan [Internet]. Journal of the American Dietetic Association. 2005;105:445–8.
Available:<http://dx.doi.org/10.1016/j.jada.2004.12.030>
 16. Sheldon P. Pressure To Be Perfect: Influences on College Students' Body Esteem [Internet]. Southern Communication Journal. 2010;75:277–98.
Available:<http://dx.doi.org/10.1080/10417940903026543>
 17. Sridharan G, Ramani P, Patankar S, Vijayaraghavan R. Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma. J Oral Pathol Med. 2019 Apr;48(4):299–306.
 18. Taras H. Nutrition and Student Performance at School [Internet]. Journal of School Health. 2005;75:199–213.
Available:<http://dx.doi.org/10.1111/j.1746-1561.2005.00025.x>
 19. Egbuna C, Mishra AP, Goyal MR. Preparation of Phytopharmaceuticals for the Management of Disorders: The

- Development of Nutraceuticals and Traditional Medicine. Academic Press. 2020;570.
20. Ezhilarasan D. Critical role of estrogen in the progression of chronic liver diseases. *Hepatobiliary Pancreat Dis Int.* 2020 Oct;19(5):429–34.
 21. Soldavini J, Berner M, Da Silva J. Rates of and characteristics associated with food insecurity differ among undergraduate and graduate students at a large public university in the Southeast United States [Internet]. *Preventive Medicine Reports.* 2019;14:100836. Available:<http://dx.doi.org/10.1016/j.pmedr.2019.100836>
 22. Lowry R, Thompson B, Lowry R. How much do the general public want to be involved in decisions on implementing water fluoridation? [Internet]. *British Dental Journal.* 2000;188:500–2. Available:<http://dx.doi.org/10.1038/sj.bdj.4800521a>
 23. Al Taki A, Khalesi M, Shagmani M, Yahia I, Al Kaddah F. Perceptions of Altered Smile Esthetics: A Comparative Evaluation in Orthodontists, Dentists, and Laypersons. *Int J Dent.* 2016 Sep 28;2016:7815274.
 24. von Bothmer MIK, Fridlund B. Gender differences in health habits and in motivation for a healthy lifestyle among Swedish university students. *Nurs Health Sci.* 2005 Jun;7(2):107–18.
 25. Ranjit N, Nielsen A, Akhavan N, Denis L, Janda K, Jovanovic C, et al. Outcomes of a Community-Wide Health Intervention in a Low-Income, Primarily Hispanic Community: The Go! Austin/Vamos! Austin (GAVA) Initiative. *Health Promot Pract.* 2020 Oct 9;1524839920961365.
 26. Gowhari Shabgah, A. et al. Shedding more light on the role of Midkine in hepatocellular carcinoma: New perspectives on diagnosis and therapy', *IUBMB Life.* 2021;73(4):659–669.
 27. J, P. C, et al. Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study', *Clinical Implant Dentistry and Related Research.* 2018;531–534. DOI: 10.1111/cid.12609
 28. Kamath SM, et al. 'In vitro augmentation of chondrogenesis by Epigallocatechin gallate in primary Human chondrocytes - Sustained release model for cartilage regeneration', *Journal of Drug Delivery Science and Technology.* 2020;101992. DOI: 10.1016/j.jddst.2020.101992.
 29. Mudigonda SK, et al. 'Non-suturing microvascular anastomosis in maxillofacial reconstruction- a comparative study', *Journal of Cranio-Maxillofacial Surgery.* 2020;48(6):599–606.
 30. Nambi G, et al. Spinal manipulation plus laser therapy versus laser therapy alone in the treatment of chronic non-specific low back pain: A randomized controlled study', *European Journal of Physical and Rehabilitation Medicine.* 2018;54(6):880–889.
 31. Prakash AKS, Devaraj E. 'Cytotoxic potentials of S. cumini methanolic seed kernel extract in human hepatoma HepG2 cells', *Environmental Toxicology.* 2019;1313–1319. DOI: 10.1002/tox.22832
 32. Rajakumari R, et al. 'Grape seed extract-soluplus dispersion and its antioxidant activity', *Drug Development and Industrial Pharmacy.* 2020;46(8):1219–1229.
 33. RH, et al. 'CYP2 C9 polymorphism among patients with oral squamous cell carcinoma and its role in altering the metabolism of benzo[a]pyrene', *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology.* 2020;306–312. DOI: 10.1016/j.oooo.2020.06.021
 34. Santhakumar P, et al. Ethanolic Extract of Capparis decidua Fruit Ameliorates Methotrexate-Induced Hepatotoxicity by Activating Nrf2/HO-1 and PPARγ Mediated Pathways', *Indian Journal of Pharmaceutical Education.* 2021;55(1s): S265–s274.
 35. Saraswathi I, et al. 'Impact of COVID-19 outbreak on the mental health status of undergraduate medical students in a COVID-19 treating medical college: a prospective longitudinal study', *Peer J.* 2020;8:e10164.

© 2021 Raj et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/78822>