

# Journal of Pharmaceutical Research International

**33(60B): 1304-1310, 2021; Article no.JPRI.80096 ISSN: 2456-9119** (Past name: British Journal of Pharmaceutical Research, Past ISSN: 2231-2919, NLM ID: 101631759)

# **Meditations on Relieving Alcohol Addiction**

# Suraj Behere <sup>a\*†</sup> and Ajay Meshram <sup>b#</sup>

<sup>a</sup> Jawaharlal Nehru Medical College, Datta Meghe Institute of Medical Sciences, Wardha, India. <sup>b</sup> Department of Biochemistry, Jawaharlal Nehru Medical College, Datta Meghe Institute of Medical Sciences, Wardha, India.

#### Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

#### Article Information

DOI: 10.9734/JPRI/2021/v33i60B34747

**Open Peer Review History:** 

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/80096

**Review Article** 

Received 15 October 2021 Accepted 20 December 2021 Published 23 December 2021

# ABSTRACT

Although prenatal alcohol causes growth retardation, mood disorders, adverse outcomes, and congenital disabilities, fetal alcohol spectrum disorders are not well diagnosed. In the world of alcohol spectrum disorders, 0.77, in Europe and North America, a high increase of 2-5%, which highlights the need for increased diagnostics and treatment. However, the history of self-reported breastfeeding has been weakened by fidelity, the absence of physical symptoms, and disorders of the fetus alcohol spectrum—There are various diagnostic differences in the system and ways to slow progress in diagnosing and controlling the disease. Neuroimaging brain structure, cortical Development, White Matter microstructure, and fetal alcohol spectrum disorders show abnormalities in functional connections in human beings. These approaches to disability development change as well as integration, high performance, memory, hearing, vision, behavior, ability to move. Promising trials of nutritional interventions and psychiatric medications continue with the goal of treatment. With 6,837 participants from the United States, the Netherlands, Italy, and Australia in 27 research projects, this book review evaluates the effectiveness of this process, which includes mindfulness, vipassana, and meditation. Meditation practitioners were able to reduce the reasons for drinking and get rid of alcohol cravings by strengthening awareness and self-control, which led to increased alcohol-related self-control, reduced stress, and eventually stopped drinking excessively. These results support the use of meditation in alcohol treatment programs. In addition, suggestions are being made to improve this strategy to effectively address

<sup>&</sup>lt;sup>†</sup>1<sup>st</sup> Year MBBS student;

<sup>&</sup>lt;sup>#</sup>Assistant professor;

<sup>\*</sup>Corresponding author: E-mail: beheresuraj3159@gmail.com;

alcohol abuse. -Meditation is a promising cure for alcoholism. This series of 16-week cases was designed to gather initial data on the effectiveness of meditation and evaluate the feasibility of research methods. Nineteen adults who relied on alcohol were enrolled in a hospital stay. Patients were randomized to either a standard course of chlordiazepoxide four times daily with additional medication as needed (fixed-schedule therapy) or to a treatment regimen that provided chlordiazepoxide only in response to the development of the signs and symptoms of alcohol withdrawal (symptom-triggered therapy). The need for administration of "as-needed" medication was determined using a validated measure of the severity of alcohol withdrawal.

Keywords: Meditations; alcohol; addiction; abuse.

#### 1. INTRODUCTION

Liquor misuse stays a real impediment for all ages, from the immature to the older, and makes physical, mental, conduct, monetary, family, and social problems within folks and over 300000 passings per annum overall prescribed with the impacts of a high danger issue connected within more than two hundred physical and mental and incapacities, sicknesses as well as malignant organ growth, diabetes, vas disappointment, mishap vessel, dementia, and crazy issues. This makes quantitative and immaterial social cash charges [1]. The most cover upbeat defrayment, lower utility, law implementation prices, defrayment on rejection and instruction, and social money plans; The last possibility incorporates inferiority of life and antagonistic family connections that cause native [2]. Additional unfortunate still, maternal liquor prompted craniate liquor condition, maybe prompts hereditary liquor abuse that hurts babies for the length of their lives. A good assortment of medicines are used to battle liquor addiction, principally with medicine, conduct alteration or joined methodologies that manufacture positive outcomes: Β. medication with narcotic antagonist, or associated with intellectual, social treatment. Contemplation, a non-noisy follow that develops the energy of medicine conduct, has in addition been used for substance misuse and backslides since the Seventies, for example, supernatural reflection: the amount that{increasingly progressively more and additional} more Buddhist structures are being settled for as well as and careful reflection. For example, Associate in Nursing ebb and flow reports the viability of care primarily based on a lapse anticipation program in 168 medication or liquor dependent members, which was upheld by alternative examination [3].

This writing audit offers an overview of the impacts of utilizing contemplation on liquor misuse utilizing twenty-seven important electronic info bases, for instance, nation Nursing Index, Medical Library, MEDLINE; the incorporation rules are four: (1) use only meditation (2) only investigate problem alcohol use, (3) apply empiric techniques, and Exclusion rules include (1) duplications and (2) Theoretical exploration, literature review, and dissertation [4].

#### 2. FINDINGS AND DISCUSSION

The twentv-seven ventures investigated concerned six, 837 members from us, Italy, European nation, and Australia, and three,816 (55.8%) teens or undergrads, nine (33.3%) of whom were trial examines. Simply three randomized controlled preliminaries (11.1%), 484 (7.1%) members were directed, and eighteen (0.3%) members took Associate in Nursing interest in another excellent subjective review [5]. Albeit the writing search was on the market to any or all kinds of reflection, the discharge recorded Buddhist-situated kinds of contemplation, basically known with care (n = 24). 88.9%) [6].

#### 2.1 Mindfulness-Based Meditation

As properties in analysis connected with fixation, the five parts of care deciphered by Western researchers embrace noticing (thoughtfulness concerning within and out of doors conditions), symbolic process (word articulation of internal encounters), acting with attentiveness (present second exercises), non-judging (non-assessment to musings and sentiments), and non-reactivity to progression inward encounters (free of contemplations and feelings) [7]. B is based on the proper octuple manner, a focal Buddhist rule, the two essential parts of care are memory and Memory centers attentiveness. around recollecting the excellencies to accomplish inward remotion through the profound perception of the body, sentiments, psyche, and peculiarities that frame the conscious of one's soiled natural impermanence. impermanence. and the

consistent emerges. These will be accomplished through careful preparation, which prompts inherent knowing characteristics like fervid responses and individual shortcomings (counting voracity, On these lines, care has been loosely taken on for physical and emotional well-being medicines in individual and authoritative settings and covers a good age vary from youngsters to the previous. Its adequacy in chronic drug use and substance-actuated dysfunctional behavior is drawing within the thought of the medical services framework [8]. Traits independence analysis incorporate the 5 parts of care deciphered by Western researchers, perception (regard for the within and outer climate), (verbal articulation portrayal of interior encounters), cognizant activity (exercises at the time present), no - analysis (no assessment of musings and sentiments) and no reactivity to inward encounters (free progression of and contemplations sentiments). Care contemplation adds to functioning on these aspects, which, for example, are connected with a negative relationship with explanations behind drinking and liquor connected to pressure. 13 members visited day by day two-hour conferences through weekday for a substantial length of your time, rehearsing half-hour of care reflection reception day by day, that showed enhancements conduct and character in development, nonetheless in addition a decrease within the danger of making a character-also, the lapse of liquor abuse. Additionally, 207 undergrads with dangerous liquor use were offered care getting ready, directed by a 10minute recording, to examine the association between thought processes in drinking, drinking conduct, and care. The outcomes showed Associate in Nursing an unfavorable relationship between conscious activity and the probability of unsafe liquor utilization attributable to less liquor utilization as a justification for adapting to diminish negative feelings. This affiliation likewise applies to the non-judgment issue. Moreover, additional cognizant activity is said with a lower likelihood of liquor ill-usage as a justification for the consistency to appease strain from others. These were thoroughbred by Sir Joshua Reynolds and O'Connor from their study of seventy-six sources. Further, the} study of 316 respondents added a reverse association among portrayal and liquor utilization [4]. A perception from an additional examination showed Associate in Nursing irrelevant association among perception and non-reactivity to inward liquor abuse [9]. encounters and Care remarkably prompts an additional grounded,

optimistic verifiable assumption and an additional vulnerable negative implicit assumption in lightweight of the very fact that it builds attentiveness and willingness and diminishes appended thinking. Such qualities result in higher leader management (more important guideline of thought and conduct) and a decrease in plan concealment that directs recuperating reactivity and deactivates liquor signs. Similarly, as more developing poise and lightening pressure programmed reactivity and impulse. as liquor elaborated eighteen immature by purchasers during a subjective report following collaborating during a 10-week care organized intercession, care debilitates healing programmed liquor inspiration nonetheless supports liquor connected self-viability. Thus, it adds to decreasing liquor utilization and dealing on emotional well-being by cashing in on an 8week reflection course joined with a day 30minute practices reception and facilitating from pressure. gloom, nervousness, and liquor connected upbeat-objective problems in life [10].

# 2.2 Vipassana Meditation

Reflection underlines the impression of fantastic reality, that is, H. the thought of the thought world, as so much as ephemerality, enduring, and impermanent self, and eventually makes progress toward liberation. As declared by 173 members WHO rehearsed contemplation for eight to ten over a 10-day course, a nonrandomized review showed an enormous reduction in undesirable musings regarding liquor use. This exploration gave positive pointers; additional dependable info will be essential to analyze the impacts of this Buddhist reflection strategy on liquor compulsion [11].

# 2.3 Transcendental Meditation

Despite the high cost of occupational stress, few studies have empirically documented effective methods for alleviating stress and promoting development. This employee three-month prospective study evaluated the effects of the Transcendental meditation (TM) technique on reduction, health, and stress employee development in two settings in the automotive industry: a large manufacturing plant of a Fortune 100 corporation and a small distribution sales company. Employees who learned TM were compared to controls similar in the worksite. position. demographic, iob and pretest characteristics [12]. Regular meditators improved significantly more than controls (with irregular meditators scoring in between) on multiple measures of stress and employee development. physiological reduced including arousal (measured by skin conductance levels) during and outside TM practice: decreased trait anxiety. job tension, insomnia, and fatigue, cigarette and hard liquor use; improved general health (and fewer health complaints); and enhanced employee effectiveness, job satisfaction, and work/personal relationships. Principal components analysis identified three factors underlying this wide range of improvements through TM: "occupational coherence." "physiological settledness," and "job and life satisfaction [13]. The "effect size" of TM in reducing skin conductance, trait anxiety, alcohol/cigarette use, and in enhancing personal development (relative to the control condition) in these business settings was substantially larger than for other forms of meditation and relaxation reported in four previous statistical metaanalyses.

# 2.4 Qigong Meditation

A temporary residential treatment program offers Qigong meditation, which includes relaxation, breathing, guided images, meditation, and mindfulness to achieve a calm state. Initially, customers preferred to volunteer to participate in Qigong meditation during the evening holidays. He then participated in meditation or stress management and relaxation training (SMART) twice a day as part of planned treatment. Weekly questionnaire The 248 participants completed four weeks to assess changes in their treatment outcomes. Participants in the meditation group were also evaluated for quality of meditation to assess the link between quality and effect of treatment. Qigong meditation seems to have a positive effect on the effects of addiction treatment, which has positive effects as a small stress management program. The results of the meditation are very encouraging. Meditation therapy may be more effective or acceptable for women who use drugs than men. Further research is needed to explore ways to improve substance abuse and attention skills [14].

# 2.5 Raja Yoga Contemplation

The review Brahmakumans Raja Yoga underlines the decreasing consequences for actual changes, for example, pulse, respiratory rate, isotolic and diastolic circulatory strain contrasted with the pre-reflection time of contemplation. Diminished physiological inconstancy demonstrates an adjustment of the

pathogenesis state vet to be determined pieces of the autonomic sensory system. with compassion incitement expected to diminish during Raja Yoga practice [15]. Thusly thoughtful incitement likewise decreases heart and weight which works on cardiovascular boundaries. Comparative discoveries, like our review, Gupta S. In 23 subjects, et al showed an abatement in thoughtful upgrades in pulserate, respiratory rate, and circulatory strain levels. It is accepted that reflection steadily diminishes the strength of sympathy which prompts a superior harmony among Tempathy and parasympathetic. This prompts a high metabolic state which brings down pulse and circulatory strain. 15 Lactic corrosive lack because of contemplation is viewed as a manifestation of a tranquil and quiet perspective. The parasympathetic cholinergic drive reaction by the vagus hub to the SA hub lessens the pulse, doesn't inspire compassion toward the theta 1, beta 2 receptors of the SA hub, or decreases the adrenaline motivation which further diminishes the pulse.

Parasympathetic cholinergic expansion of the conduits diminishes fringe opposition prompting a lessening in diastolic pulse, a reduction in thoughtful action decreases the constrictor impact brought about by flour gasopters which thus brings down the systolic circulatory strain Basimerfectic clinical responses that cause a negative topic activity to decrease cardiovascular yield, diminishing heart withdrawal, and along these lines bringing down systolic pulse. Diminished thoughtful reaction lessens withdrawals. The reduction in respiratory rate after reflection is manifested by a dynamic expansion in vaginal boosts towards the respiratory focus. Notwithstanding actual measures, our review checked out the emotional well-being of those subjects Focuses on the helpful impacts of reflection on the people who practice Raj Yoga contemplation, like controlling displeasure, to Raj Yoga. What's more, Raja Yoga can help intellectually upsetting subjects. 14,16 The consequences of different examinations additionally affirm the flow brings about an exploration study, the clinical part of Raja Yoga Education and Research Foundation analyzed 189 individuals who were dependent on tobacco prior to beginning the act of Raj Yoga. In 181.74% of individuals quit smoking in one month Another 19 became dependent in only 3 months. Along these lines, in only 3 months, 931 individuals became dependent on drugs. This implies Rajyog to dispose of medications [16].

Mental state. After an extensive stretch of learning and rehearsing state voga contemplation. negative/inefficient stress. reasoning and crabbiness just as stopping smoking, liquor and smoking appears to work on the legs of reflection. Zero in on passionate movement identified with feeling as reflection has been found in many examinations to decrease uneasiness and increment its constructive outcomes, also [17].

# 3. CONCLUSION

It is equally protected, reasonable, non-clinical, and non-intrusive. The positive signs from the inspected show the impacts of contemplation on liquor use problems as so much as a good scope of outcomes from a healing and preventive perspective, as well as diminished inspiration to drink, higher conveyance of liquor and distended confidence known with liquor. This coaching likewise helps liquor connected psychological well-being problems, like discouragement and nervousness. Whereas it's a probable proportion of neurotic liquor habit and lapse, all the additional all around planned precise examination is predicted to assemble considerable and solid proof.

#### 4. RECOMMENDATIONS

Advocated throughout the Fifties and created from Buddhist thoughts, supernatural contemplation could be a deliberate profound respiration procedure, settled to mantras, to develop internal joy corresponding to harmony and equilibrium. A vital study of one, 862 lecturers WHO follow supernatural contemplation discovered that this coaching may decrease utilization what is more, and also the cluster itemized the implications of their task in this fourhundredth of 216 members WHO rehearsed this reflection for a protracted time guit drinking and hour guit drinking within 2539 months. Even so these OK outcomes. late scholastic exploration on this method appears to possess become dormant with regards to liquor misuse [18].

# 5. LIMITS AND FUTURE

Exploration Directions analysis techniques and strategies of will be more to look at their accessibility against liquor misuse. Within the initial place, simply thirty third of the analyzed zeroed in on mediation programs, whereas the lay trusted phone or net overviews. Distended trial analysis is enticing to reinforce the promising info on however reflection will address hard-core

drinking. Second. 3 randomised controlled preliminaries didn't produce enough logical outcomes [19]. A randomised controlled exploration configuration is required to collect solid informational indexes during a proof primarily based manner that limits analysis inclination. Third, within the explored papers a cross-sectional set up while not follow-up was used, that makes it laborious to make a decision its drawn out viability [20]. During this manner, longitudinal investigations ar to expand the applications to liquor connected problems. Fourth. hiah to bottom conferences. notwithstanding whether or not in a private or social scene, will provides a additional profound arrangement and purpose by purpose record of the lived encounters of people with liquor use issues. Consequently, subjective examination could be a recommended strategy. At last, broadening kinds of contemplation, like Taoist reflection, is relied upon to increase the alternatives for each upbeat and alcoholics [21-29].

# SUGGESTIONS

Normal attributes of addictive conduct ar avid weakness, incitement, and loss of management. which may be improved by contemplation, for example, a decrease in wanting for liquor. The implications of this survey support the extended uses of reflection to compulsion treatments, for instance, card-playing, undue utilization of online media and sexual fixation. Nevertheless, having a licensed reflection professional is key to remain faraway from dreams, daydreams, or pipedreams whereas rehearsing. This can be the place wherever Associate in Nursing overall rule becomes Associate in Nursing integral factor: follow, additional proficient strategies, and additional distinguished viability. Consequently, perform contemplation suggested iť's to systematically as Associate in Nursing adapting methodology and to coordinate it into exercises of day by day living: B. Strolling, indestion and preparation. This shows paying outcomes for end of the day remedial and preventive functions. Wellspring of subsidizing Support Preparation of this text didn't get any funding. Statement of interest There is no unreconcilable scenario to pronounce regarding this composition submitted for distribution.

# NOTE

The study highlights the efficacy of "Raja Yoga, Vipassana" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

#### DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

# CONSENT

It is not applicable.

# ETHICAL APPROVAL

It is not applicable.

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

# REFERENCES

- 1. Patton R, Deluca P, Kaner E, Newbury-Birch D, Phillips T, et al. Alcohol Screening and brief intervention for adolescents: the how, what and where of reducing Alcohol consumption and related harm among young people. Alcohol Alcohol. 2014;49: 207-212.
- Chikere EIC, Mayowa MO. Prevalence and perceived health effect of alcoholUse among male undergraduate students in Owerri, South-East Nigeria: A descriptive Cross-sectional study. BMC Public Health. 2011;11(1:18): 1-6.
- 3. Hanes M. Effects and consequences of underage drinking. Juvenile Justice Bulletin. 2012;1-11.
- 4. Rogers J, Wiese B. Geriatric drinkers: Evaluation and treatment for alcohol Overuse. BC Medical Journal. 2011;53(7): 353-356.
- 5. World Health Organisation, Alcohol;2015.
- Shield KD, Parry C, Rehm J.Chronic diseases and conditions related to alcohol Use. Alcohol Res. 2013;35:155-173. [Crossref]

- Single E, Collins D, Easton B, Harwood H, Lapsley H, et al. International Guidelines for estimating the costs of substance abuse. 2<sup>nd</sup> ed. Geneva, Switzerland: World Health Organisation;2003.
- 8. Burke TR1. The economic impact of alcohol abuse and alcoholism. Public Health Rep. 1988;103:564-568. [Crossref]
- 9. Reinaldo AM, Pillon SC. Alcohol effects on family relations: a case study. Rev Lat Am Enfermagem. 2008;16:529-534. [Crossref]
- 10. World Health Organisation. Global status report on alcohol 2004. Geneva, Switzerland: World Health Organisation; 2004.
- 11. Saitz R, Mayo-Smith MF, Roberts MS, Redmond HA, Bernard DR, Calkins DR. Individualized treatment for alcohol withdrawal. A randomized double-blind controlled trial. JAMA. 1994;272(7):519-23. PMID: 8046805.
- Abrams AI, Siegel LM. The Transcendental Meditation program and rehabilitation at Fol-som State Prison: A cross-validation study. Crim. Justice Behav. 1978;5:3–20 [Crossref], [Web of Science ®], [Google Scholar]
- Alexander CN. Ego development, personality and behavioral change in inmates practicing the Transcendental Meditation technique or participating in other programs: A cross-sectional and longitudinal study. Dissert. Abstr. Int. 1982;43(2)539-B [Google Scholar]
- 14. Alexander CN, Davies JL, Dixon CA, Dillbeck MC, Druker SM, Oetzel RM, Muehlman JM, Orme-Johnson DW. Growth of higher stages of consciousness: psychology The Vedic of human development. Higher stages of human development: Perspectives on adult growth, C. N. Alexander, EJ Langer. Oxford University Press, New York;1990 [Google Scholar]
- Alexander CN, Langer E, Newman R, Chandler H, Davies JL. Transcendental Meditation, mindfulness and longevity: An experimental study with the elderly. J. Pers. Soc. Psychol. 1989;57:950–964 [Crossref], [PubMed], [Web of Science ®], [Google Scholar]
- Alexander CN, Rainforth MV, Gelderloos P. Transcendental Meditation, selfactualization, and psychological health: A conceptual overview and statistical metaanalysis. Journal of Social Behavior and

Personality. 1991; 6(5)189–247 [Google Scholar]

- Aron EN, Aron A. Transcendental Meditation program and marital adjustment. Psychol. Rep. 1982;51:887– 890 [Google Scholar]
- Ballou D. The Transcendental Meditation program at Stillwater prison. Orme-Johnson and Farrow;1977 [Google Scholar]
- Bangert-Drowns RL. The effects of schoolbased substance abuse education: A meta-analysis. J. Drug Educ. 1988; 18:243–264
- 20. Benson H, Wallace RK. Decreased drug abuse with Transcendental Meditation: A study of 1, 862 subjects. Drug abuse: Proceedings of the international conference, C. J.D. Zarafonetis. Lea and Febiger, Philadelphia;1972 [Google Scholar]
- Berg WP, Van Den Mulder B. Psychological research on the effects of the Transcendental Meditation technique on a number of personality variables. Gedrag: Tijdschrift voor Psychologic 1976;4:206–218 [Google Scholar]
- Behere, Prakash B., Aniruddh P. Behere, and Debolina Chowdhury. "Rise in Cases of Alcohol Withdrawal Following Lockdown in Dry District of Wardha. Journal of Neurosciences in Rural Practice. 2020;11(03):478–80. Available:https://doi.org/10.1055/s-0040-1712554.
- 23. Ghogare, Ajinkya Sureshrao, Pradeep Shriram Patil, and Ganpatlal Kodarbhai Vankar. "A Protocol of Case Control Study of Childhood Trauma and Alexithymia in Persons with Alcohol Dependence Syndrome. Journal of Clinical and Diagnostic Research. 2020;14(9).

Available:https://doi.org/10.7860/JCDR/20 20/44593.14018.

 Gedam, Sachin Ratan, Ajab Dhabarde, Pradeep S. Patil, Animesh Sharma, Kanika Kumar, and Vijay Babar. "Psychiatric Comorbidity, Severity of Dependence and Liver Enzymes Dysfunction among Alcohol Dependent Individuals: A Cross-Sectional Study from Central Rural India." Journal of Clinical and Diagnostic Research. 2019;13(4):VC1–5. Available: https://doi.org/10.7860/.ICDB/20

Available:https://doi.org/10.7860/JCDR/20 19/40368.12759.

- 25. Ghogare, Ajinkya Sureshrao. Study of Cognitive Impairment in Patients of Alcohol Use Disorder. Indian Journal of Psychiatry. 2019;61(9)3:S583–84.
- 26. Joshi, Amey, Lipsy Modi, and Imran Ali Shivji. Comparison of Total Self-Stigma between Schizophrenia and Alcohol Dependence Patients. Indian Journal of Psychiatry. 2018;60(5):125–26.
- Patel, Tejas, V, Mehul J. Brahmbhatt, and Ganpat K. Vankar. "Prevalence of Alcohol Use Disorders in Hospitalised Male Patients. Archives of Psychiatry and Psychotherapy. 2018;20(4):47–55. Available:https://doi.org/10.12740/APP/991 47.
- Patel D, Gandhi Z, Desai R, Raina J, Itare V, Haque FA, et al. Impact of alcohol use disorder on stroke risk in geriatric patients with prediabetes: A nationwide analysis. International Journal of Clinical Practice.
- 29. Bhainsora RS, Patil PS, Ghogare AS, Vankar GK. A cross-sectional study of prevalence and types of sexual dysfunction among married male patients with alcohol dependence syndrome attending tertiary healthcare center from Central Rural India. Journal of Education and Health Promotion. 2021;10(1).

© 2021 Behere and Meshram; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

> Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/80096