



Ellis Van Creveld Syndrome: A Case Report

**Sarath Kumar^{a*‡}, S. Padmavathi^{a‡}, G. Anuradha^{a‡}, A. Kannan^{a‡}
and C. L. Krithika^{a‡}**

^a Department of Oral Medicine and Radiology, SRM Dental College, Ramapuram, SRM University, Chennai, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2021/v33i61B35526

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/83240>

Case Report

Received 20 October 2021

Accepted 25 December 2021

Published 28 December 2021

ABSTRACT

Ellis Van Creveld Syndrome (EVC) is a infrequent inborn genetic disorder with an x-linked recessive pattern of inheritance. This is recognized by bilateral accessory little finger in upper extremity, short limbs, ectodermal dysplasia affecting teeth and nails, and congenital cardiac defect. The overall prevalence of this disease is 7 in 1000000. In this current case, a 9year old female patient reported with typical general and oral manifestations, which were key diagnostic features of EVC syndrome.

Keywords: *Ellis van creveld syndrome; congenital heart defect; autosomal recessive.*

1. INTRODUCTION

EVC is a mesenchymal ectodermal dysplasia first reported by two Pediatricians named Richard W.B Ellis and Simon van Creveld in 1940 [1]. It is a infrequent congenital x-linked recessive

syndrome due to a genetic abnormality on chromosome 4p16 [2]. It is brought by a mutation in EVC1 and EVC2 genes with parental consanguinity in 30% of cases [3,4,5]. In 1964, Mukusick et al reported many affected individuals in the Amish community of Lancaster Country,

[‡]Post-Graduate Student;

[#]Professor and Head;

[‡]Professor;

[†]Reader;

^{*}Corresponding author;

Pennsylvania of USA and in western part of Australia. The frequency of this disorder in the Amish population is 1 in 5000 actual births, whereas in the other communities is 7 in 1000000 [4,5].

EVC Syndrome presents with the following tetrad of clinical manifestations [6] which includes; First most typical hallmark triad is skeletal dysplasia of the os longum, which results in a relatively small stature and elongated torso, resulting in a hazardous ossification defect [5]. The intensity of short limbs rises as they progress from proximal to distal. Lateral ray polydactyl of upper extremities with a extra digit usually begins on the side of little finger. With large hands and feet, sausage-shaped fingers are common. Cloutson syndrome with dystrophy of nails, weak and scanty hair, and oral manifestation [7]. In 50-60% of cases, there is inborn heart illness, the most prevalent of which involves only one atrium chamber and ventricular septal defect [5].

2. CASE REPORT

A 9-year-old young female patient who reported to the Department of Oral Medicine and Radiology with a one-year history of disparity among her upper and lower front teeth. Exfoliation of primary dentition and lack of permanent teeth eruption in the upper anterior region was also mentioned by the patient. Medical history revealed congenital heart

disease (Arterio ventricular canal defect) for which she had undergone cardiac surgery at the age of 1 year. There is no history of the parents marrying consanguineously and family history is also noncontributory.

The child had a small stature and both limbs are shortened and malformed on general inspection. (Fig.1). She also presented with strabismus (Fig. 2), cubitus valgus, narrow chest, hypoplastic fingernails, the big toe and the other toes have a significant disparity between them in the right foot (Fig. 4). The patient cognitive capability was within a reasonable range. The intraoral evaluation divulged morphological alterations in upper and lower anterior, midline diastema, highly attached labial frenum in the maxillary anterior region, absence of both maxillary lateral incisors (Fig. 3).

Panoramic radiograph confirmed the absence of maxillary laterals (Fig. 5A). Shortening of the extremities, genu valgum, has been seen on a radiograph of the upper and lower limbs (knock knees), bilateral postaxial polydactyly, hand wrist radiograph showed the hamate and capitate bones, as well as the 5th and 6th metacarpal bones, are fused together (Fig. 5B). On the basis of clinical and radiological evidence with an associated congenital heart defect, the patient was provisionally diagnosed with Ellis van Creveld syndrome.



Fig. 1. Patient with short stature measuring about 119 cm in height



Fig. 2. Patient show strabismus, on clinical examination



Fig. 3. On intraoral examination, conically shaped tooth in upper and lower anterior. The frenal fibers stretch upto the palatine papilla, crossing the alveolar process



Fig. 4. Extra digit is evident on both upper and lower limbs with hypoplastic nails and the big toe is separated from the other digits by a significant distance



Fig. 5A and B. Orthopantogram confirms conical-shaped maxillary and lower front teeth with congenitally missing maxillary laterals. The hand-wrist X- ray depicts union of carpals and extra digit on side of little finger.

3. DISCUSSION

Chondroectodermal dysplasia is a infrequent x-linked recessive disorder. When enhanced nuchal translucency is observed on ultrasound, the detection of this disease can be confirmed as soon as the 18th week of pregnancy [4]. The prevailing clinical sign is acromesomelic dwarfism owing to defect in ossification affecting

tubular bones, which emanates in the shortening of bones of limbs [8]. Other attributes consists of polydactyl usually bilateral accessory little finger which is commonly present in upper limbs and on the ulnar side of limbs and only in 10% of reports, it can be present in lower limbs [3]. EVC individuals also had cubitus valgus, genu valgum, dysplastic fingers and nails, and narrow chest [4]. Inborn cardiac defects are present

among 60% of cases which include defects of bicuspid and right atrioventricular valves, ductus arteriosus closure, atrial and ventricular septal defect which are leading reasons for reduced life expectancy [9]. Some inconstant clinical features of EVC are congenital cataracts, nephrocalcinosis, strabismus, retinitis pigmentosa, central nervous anomalies, hypoplastic penis, and hematological anomalies [10].

The characteristic oral findings of this syndrome which aids in diagnosis which are, union of upper lip to the gingival margin, leading to loss of mucobuccal fold, partial non union of upper lip, multiple small accessory frenum, tongue tie, malignment of teeth, atypically small teeth, missing of lower successor anterior teeth and enamel hypoplasia [11]. In our present case, most of the clinical signs and symptoms such as acromesomelic dwarfism, bilateral postaxial polydactyl, narrow chest, cubitus valgus, hypoplastic fingers and nails, genu valgum, and previously treated congenital heart disease were seen. In addition to this, some inconstant features such as strabismus and unilateral postaxial polydactyl in the left foot and anodontia of maxillary laterals are not reported in much of the literature. The definitive diagnosis of this syndrome is genetic-based where homozygosity for mutation in EVC1 and EVC2 genes which is detected by direct sequencing. However, the genetic mutation is positive only in 2/3rd of patients [12]. Due to less availability of genetic studies, the diagnosis of the current case was arrived by clinical and radiological findings. Although inborn cardiac defect and breathing illnesses are the leading causes of death in this disorder, those who sustain oneself childhood will get a standard survival rate. The individual who survived the oldest was eighty two years old [13].

Differential diagnosis of this case was curly hill syndrome, Jeune syndrome, orofacial digital syndrome, Morquio syndrome, and achondroplasia. Since there is no cure for EVC [14], Management focuses on symptomatic relief followed by a multidisciplinary approach. Some studies have found somatotropin insufficiency in this disorder, along with supplementation of somatotropin has been shown to improve these patients' stature [15].

4. CONCLUSION

EVC Syndrome is a occasional congenital genetic disorder which needs a multidisciplinary

approach for a suitable diagnosis and management. Patients with this syndrome encounter a high mortality index because of cardiovascular and respiratory complications. Hence dentists play an essential role in early detection and establishing management protocols. Dentists can help improve the esthetics and function, overcoming psychological trauma, and enhance the wellbeing of patient.

CONSENT

The authors attest to having gotten all necessary patient consent papers. The patient(s) mother has/have consented in the form for her daughter's photos and other diagnostic evidence to be published in the paper. The consent individual mother is aware that their identities and surnames will be not be advertised, and that while every effort will be taken to keep their identities hidden, privacy could not be guaranteed.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Muensterer OJ, Berdon W, McManus C, Oestreich A, Lachman RS, Cohen MM Jr, Done S. Ellis-van Creveld syndrome: its history. *Pediatr Radiol.* 2013 Aug;43(8):1030-6.
2. Howard TD, Guttmacher Ae, McKinnon W, Sharma M, Mckusick VA, Jabs EW. Autosomal dominant postaxial polydactyly, nail dystrophy, and dental abnormalities map to chromosome 4p16, in the region containing the Ellis-van Creveld syndrome locus. *Am. J. Hum. Genet.* 1997;61:1405-1412.
3. Veena KM, Jagadishchandra H, Rao PK, Chatra L. Ellis-Van Creveld syndrome in an Indian child: a case report. *Imaging Sci Dent.* 2011;41:167-170.
4. Hegde K, Puthran RM, Nair G, Nair PP. Ellis-Van Creveld syndrome report of two siblings. *Br Med J Case Reports.* 2011;1-8.
5. Cahuana A, Palma C, Gonzales W, Ge an E. Oral manifestations in Ellis-van creveld

- syndrome. Rep Five Cases *Pediatr Dent*. 2004;26:277-282.
6. Hattab FN, Yassin OM, Sasa IS. Oral manifestations of Ellis-van Creveld syndrome. Report of 2 siblings with unusual dental anomalies, *Journal of Clinical Pediatric Dentistry*. 1998;22:159–165.
 7. Zangwill KM, Boal DK, Ladda RL, Opitz JM, Reynolds JF, Dandy-Walker malformation in Ellis-van Creveld syndrome, *American Journal of Medical Genetics*. 1998;31(1):123–129.
 8. Gopal G, Belavadi GB. Case report of a child with Ellis-van Creveld syndrome. *Int J Pharm Biomed Res*. 2014;5:14-7.
 9. Cesur Y, Yuca SA, Uner A, Yuca K, Arslan D. Ellis-Van Creveld syndrome. *Eur J Gen Med*. 2008;5:187-190.
 10. Das D, Das G, Mahapatra TK, Biswas J. Ellis van Creveld syndrome with the unusual association of essential infantile esotropia. *Oman J Ophthalmol*. 2010 Jan;3(1):23-5.
 11. Himelhoch DA, Mostofi R. Oral abnormalities in the Ellis-van Creveld syndrome: Case report. *Pediatr Dent*. 1988;10:309-313.
 12. Ruiz-Perez VL, Tompson SW, Blair HJ, Espinoza-Valdez C, Lapunzina P, Silva EO, et al. Mutations in two nonhomologous genes in a head-to-head configuration cause Ellis-van Creveld syndrome. *Am J Hum Genet*. 2003; 72:728-32.
 13. Gorlin RJ, Cohen MM, Hennekam RCM. Syndromes affecting bone. Chondrodysplasias and chondrodystrophies. *Syndromes of the head and neck*. Oxford University Press. 2001;4:239–42.
 14. Kamal R, Dahiya P, Kaur S, Bhardwaj R, Chaudhary K. Ellis-van Creveld syndrome: A rare clinical entity, *Journal of Oral and Maxillofacial Pathology*. 2013;17(1):132–135.
 15. Versteegh FG, Buma SA, Costin G, de Jong WC, Hennekam RC, EvC Working Party. Growth hormone analysis and treatment in Ellis-van Creveld syndrome. *Am J Med Genet A*. 2007; 143:2113–21.

© 2021 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/83240>